

REVIEW ARTICLE

CONSIDERING THE IMPORTANCE OF MEDICINAL PLANTS AND NATURAL PRODUCTS AND THEIR MECHANISMS OF ACTION FOR TREATMENT OF GASTROINTESTINAL DISORDERS

Mohamad Hesam Shahrajabian, Wenli Sun

National Key Laboratory of Agricultural Microbiology, Biotechnology Research Institute, Chinese Academy of Agricultural Sciences, Beijing 100086, China

Received 25th August 2025.

Accepted 18th November 2025.

Published 1st September 2026.

Summary

Background and Purpose: The application of medicinal plants and herbs to cure diverse animal and human ailments predates recorded history. Gastrointestinal disorders have a high prevalence in human societies.

Methods: This research examined the scientific literature from 1990 to June 2025 by conducting a bibliometric analysis of the literature published on the Web of Science database, including more than one thousand articles. The information provided was obtained from randomized control experiments, analytical observations, review articles, and studies which were gathered from different literature sources such as Google Scholar, Scopus, Science Direct, and PubMed.

Results: The gastrointestinal tract is one of the most important organs in the human body, and it is vulnerable to great diversity of diseases such as infectious and parasitic disorders, constipation, gastroenteritis, reflux, diarrhea, and bloating. Some of the most important medicinal plants which have been used in different traditional medicinal sciences for the treatment of dyspepsia symptoms are chamomile (*Matricaria recutita*), cinnamon (*Cinnamomum verum*), ginger (*Zingiber officinalis*), licorice (*Glycyrrhiza glabra*), peppermint (*Mentha piperita*), and guava (*Psidium guajaba*). The most notable herbs used to treat bloating are aniseed (*Pimpinella anisum*), cassia (*Cinnamomum aromaticum*), celery (*Apium graveolens*), cinnamon (*Cinnamomum verum*), dill (*Anethum graveolens*), fennel (*Foeniculum vulgare*), ginger (*Zingiber officinalis*), parsley (*Petroselinum crispus*), pennyroyal (*Mentha pulegium*), peppermint (*Mentha spicata*), star anise (*Illicium verum*), and thyme (*Thymus vulgaris*). Phytochemicals such as seaweed-derived polysaccharide, phytohemagglutinin, and allicin have been reported as important components to induce precocious gut maturation. Phytochemicals can also change the composition of gut microbiota, digest normally indigestible molecules into bioavailable molecules, and enhance nutrient availability. *Ganoderma lucidum*, Gegen Qinlian decoction, Huang-qin decoction, and *Glycyrrhiza uralensis* Fisch. are important traditional Chinese medicine with significant effects on intestinal flora and different microbial communities.

Conclusions: The review article aims to study and survey the importance of medicinal plants and herbs while considering their mechanisms of action for treatment of gastrointestinal disorders.

Key words: Celiac disease; Gallstones; Gastritis; Gastrointestinal disorder; Medicinal plant; Peptic ulcer disease

Highlights:

1. The gastrointestinal tract is one of the main organs of the human body, which is vulnerable to various diseases.
2. Some Chemical drugs have low effectiveness with different negative effects.
3. Medicinal plants and herbs are the best alternative options which can influence common gastrointestinal diseases and disorders.
4. Medicinal herbs and plants contain high bioactive components with positive influence for the treatment of gastrointestinal disorders.
5. Medicinal plants can significantly use for gastrointestinal diseases as these diseases are the complex illnesses of different parts of the gastrointestinal tract, which can influence small intestine, stomach, esophagus, rectum, and large intestine.
6. Medicinal plants which can improve gastrointestinal health are *Matricaria chamomilla*, *Mentha spicata*, *Melissa officinalis*, *Artemisia ludoviciana*, *Flourensia cernua*, *Phoradendron californicum*, and *Turnera diffusa*.
7. Medicinal plants with natural components can provide effective and safe alternatives to synthetic medications with lower side effects and broader potential for treatment customization.

Introduction and background

Plant-based treatments and therapies are one of the most important choices for rural societies in different remote regions, especially in developing and under-developed countries (1-5). They can be used in the treatment of different infectious wounds because of their minimal toxic side effects, high effectiveness, renewable nature, and wide availability (6-10). Gastrointestinal disorders can be structural or functional (11,12), and both types may cause severe impairment of psychological wellbeing and negatively influence life quality as well as shortened life expectancy (13-16). Many non-psychoactive cannabinoids, like cannabidiol are appropriate to treat various diseases and disorders of the gastrointestinal tract (17,18). Other signs of gastrointestinal disorders are changed central nervous system processing, visceral hypersensitivity, motility disturbance, changes concerning gut microbiota, and alteration in immune and mucosal functions (19-21).

Celastrol, a triterpene obtained from the root of *Thunder of God Vine* which is very common in traditional Chinese medicine could down-regulate biglycan protein, which is essential for gastric cancer invasion and immigration, activated receptor-interacting protein 1 and 3 (RIP1 and RIP3), and finally promoted the translation of mixed-lineage kinase domain-like (MLKL) from cytoplasm to plasma membrane, inducing to necroptosis of gastric cancer cell (22). List of medicinal plants with anti-ulcer activity are *Terminalia pallida* (Terpenoids, tannins, glycosides, flavonoids, alkaloids), *Terminalia chebula* (Carbohydrates, flavonoids, alkaloids, phenols, terpenoids, tannins), *Polyalthia longifolia* (Terpenoids, alkaloids), *Spondias mombin* (Saponins, flavonoids, tannins, phenolics), *Ficus exasperate* (Tannins, flavonoids, saponins, alkaloids), *Toona ciliata* (Sterols, quercetin, terpenoids), *Calligonum cosmosum* (Flavonoids, phenols), *Tephrosia purpurea* (Flavonoids), *Carlina acanthifolia* (Essential oils, flavonoids, inulin), *Ocimum sanctum* (Eugenol, euginal, urosolic acid), *Momordica charantia* (Sterols, steroids, alkaloids, flavonoids), *Ziziphus jujuba* (Tannins, polyphenols, sterols, alkaloids, flavonoids), *Scutia buxifolia* (Tannins, polyphenols, alkaloids, flavonoids), *Madhuca indica* (Flavonoids, triterpenoids), *Maytenus robusta* (Flavonoids, steroids, triterpenes), *Casearia sylvestris* (Triterpenes, tannins), *Uleria salicifolia* (Tannins, saponins, steroids), *Asparagus racemosus* (Polysaccharides, saponins), *Kaempferia parviflora* (Flavonoids, anthrones, alkaloids), *Passiflora foetida* (Flavonoids, phenolic compounds, alkaloids, proteins, phenols), *Alhagi maurorum* (Sterols, fatty acids, flavonoids), *Bahunia purpurea* (Tannins, sterols, flavonoids), *Cressa cretica* (Proteins, tannins, glycosides, alkaloids), and *Citrullus lanatus* (Antraquinones, saponins, steroids, terpenoids, alkaloids, tannins, flavonoids, phenols) (23-29). Ethnopharmacological researches have indicated that the application of traditional medicine for gastrointestinal disease is commonly practiced in different cultures in various part of the world. Many people in different parts of the world use medicinal plants as their first choice of health care to prevent and cure of different types of gastrointestinal disorders. The main goal of this review article is to study the effects of medicinal plants and herbs by considering different mechanisms for treatment of gastrointestinal disorders.

Gastrointestinal disorders and medicinal plants

The human gastrointestinal microbiota is composed of microbial eukaryotes (protozoan and fungi), archaea, viruses and bacteria (30), although the most diversified and abundant group is bacteria (31). Mechanism of action of plant derived substances against a broad range of gastrointestinal diseases are proanthocyanidin in *Vaccinium macrocarpon* L. which can modify biofilm formation of *Enterococcus faecalis*, and *E. coli* (32-36), zingerone of *Zingiber officinale* Rosc. which can reduce heat-labile enterotoxin (LT)-induced diarrhoea in enterotoxigenic (ETEC) via blocking the binding to GM1 ganglioside receptors of ETEC (37), epigallocatechin of *Camellia sinensis* L. which can inhibit extracellular release of Vero toxin from enterohemorrhagic *Escherichia coli* O157:H7 in enterohaemorrhagic (EHEC) (38), quercetin of *Allium cepa* L. which has role in inhibition of ATPase activity, elevates extracellular phosphatase and galactosidase of *E. coli* (39,40), berberine of *Berberis vulgaris* L. which may target proteins responsible for upholding the structure of cells and for cell division of *E. coli*, *Salmonella* spp., *V. Cholerae* (41,42), eugenol of *Syzygium aromaticum* L. which can inhibit the toxin production of EHEC (43), cinnamaldehyde of *Cinnamomum verum* J. Presl which can inhibit the toxin genes and host receptor expression and reduce toxin-mediated pathology of *C. difficile* and EHEC (43,44), baicalin (5,6,7-trihydroxyflavone) of *Scutellaria baicalensis* Georgi. which can inhibit the toxin production of *H. pylori*, and thymol of *Thymus vulgaris* L. which may disturbance of the cell membrane and cytoplasm of *C. difficile* (45,46). Kemal *et al.* (47) noted that *Artemisia herba-alba* Asso, *Artemisia campestris* L., *Mentha pulegium* L., *Juniperus oxycedrus* L., *Juniperus phoenicea*, and *Lavandula angustifolia* Mill. which have been used widely in Algerian medicinal science, contain high cytotoxic and antioxidant activities as well as caffeic acid, rutin, and catechins that can stimulate gastrointestinal digestion phases.

Herbal-derived products for treating gastrointestinal infections are curry, pomegranate, and turmeric for diarrhoea (48), *Adansonia digitata* fruit for diarrhoea (49), berberine, sulfadimidine and neomycin for acute infective diarrhoea (50), berberine for antibiotic-associated diarrhoea (51), berberine and its combinations with streptomycin, chloramphenicol, sulphaimidine, and iodochlorhydroxyquinoline for acute gastroenteritis and dysentery (52), bismuth and subsalicylate for acute nonspecific diarrhoea (53). *Artemisia maritima* L. can be used for diarrhea, intestinal worms, stomachic and vomiting (54,55), *Tanacetum falconeri* Hook. Can be applied for treatment of diarrhea, vomiting and stomachic (56), *Mentha piperita* L. can be used for treatment of vomiting and diarrhea (57,58), *Hylotelephium telephioides* (Ledeb.) can be consumed for constipation (59), both *Viola serpens* Wall. and *Oxyria digyna* (L.) can be applied for treatment of stomachic, constipation, and diarrhea (60,61), *Persicaria amplexicalus* for stomach disorders (62), *Saussurea gossypiphora* D. for constipation (63), *Aconitum heterophyllum* Wall. for intestinal worms and stomachic (64), *Echinops echinatus* Roxb. For treatment of abdominal pain (65), *Rumex hastatus* D. for stomachic and flatulence (66), *Pistacia khinjuk* stocks for indigestion (67), *Plantago himalaica* Pilg. for diarrhea and stomachic (67), *Vitis vinifera* L. for intestinal worms and constipation (68), *Cicer microphyllum* Benth. for indigestion and vomiting (69), *Capparis spinosa* L. for piles and digestive (70,71), *Artemisia annua* L. for vomiting and diarrhea (72), and *Rheum australe* D. for treatment of constipation and stomachic (73,74). Phromnoi *et al.* (75) reported that *Pisidium guajava* L., and *Punica granatum* L. have high antioxidant and anti-inflammatory activities as well as notable polyphenols with considerable potential for usage in gastric ulcer therapy. Direito *et al.* (76) found that the therapeutic effects of medicinal plants such as *Piper nigrum*, *Artemisia anomala*, *Litsea cubeba*, *Agrimonia Pilosa*, and *Morus macroura* are reductions in tumor necrosis factor- α , reductions IL-1 β , interferon- γ , and caspase levels, and increased interleukin (IL)-4, IL10, and expression of antioxidant enzymes, as well as regulation of gut microbiota. *Garcinia cowa* leaf extract contains a new polyprenylated benzophenone, which is called chamuangone, and it possesses significant antibacterial activity against gastrointestinal bacteria (77).

The most common medicinal plants which are common in Persian medicinal sciences to treatment of vomiting and nausea are the roots of *Justicia adhatoda*, bulb of *Allium cepa*, peduncle of *Mangifera indica*, gall of *Pistacia intigeriima*, resin of *Pistacia lentiscus*, kernel and peel of *Pistacia vera*, leaf and seed of *Rhus coriaria*, seed and leaf of *Anethum graveolens*, leaf and root of *Apium graveolens*, seed of *Carum copiticum*, fruit of *Coriandrum sativum*, seed of *Cuminum cyminum*, root and seed of *Foeniculum vulgare*, aerial parts of *Artemisia dracunculus*, root and leaves of *Onopordum acanthium*, leaf of *Cichorium intybus*, fruit of *Berberis asiatica*, fruit of *Berberis vulgaris*, the whole plant of *Parietaria officinalis*, seed of *Descurainia Sophia*, seed of *Cannabis sativa*, the whole plant of *Humulus lupulus*, fruit and latex of *Carica papaya*, oleo-resin of *Cistus ladaniferus*, the whole plant of *Terminalia*

chebula, fruit of *Cycas revoluta*, bark and roots of *Cyperus longus*, fruit of *Elaeagnus angustifolia*, fruit of *Quercus ilex*, flower of *Hypericum perforatum*, seed and aerial part of *Thymus serpyllum*, aerial parts of *Mentha aquatica*, aerial parts of *Ocimum basilicum*, peduncle and leaf of *Mentha piperita*, seed and flower of *Teucrium montanum*, leaves of *Satureja hortensis*, root bark of *Sassafras albidum*, fruits of *Tamarindus indica* and *Trigonella corniculata*, flowers of *Punica granatum*, fruit of *Adansonia digitata*, flower of *Melia azedarach*, fruit of *Morus alba*, the whole plant of *Rheum ribes*, seeds of *Portulaca oleracea*, seed of *Nigella sativa*, root of *Thalictrum foliolosum*, leaf of *Zizyphus sativa*, fruit of *Crataegus azarolus*, fruit of *Cerasus vulgaris*, fruit of *Cydonia vulgaris*, the whole plant of *Pyrus communis*, fruit of *Prunus cerasifera*, petal of *Rosa canina*, fruit of *Citrus limonum*, seed and peel of *Citrus sinensis*, seed and fruit of *Aegle marmelos*, fruits of *Sapindus trifoliatus* and *Solanum melongena*, Rhizome of *Valeriana dioscorides*, roots of *Valeriana jatamansi*, root of *Zingiber zerumbet*, fruit of *Vitis vinifera*, and seeds of *Elettaria cardamomum* (78-83).

In the Eastern region of Saudi arabia, red tea, chamomile, thyme, senna, fenugreek, clove and myrrh used for treatment of diarrhea; fennel, thyme, chamomile, clover and fenugreek to treat constipation; chamomile, anise, mint, myrrh, and marjoram for treatment of irritable bowel syndrome symptoms; ginger, mint, fenugreek, fennel and chamomile to treat dyspepsia and indigestion; and anise, fennel, thyme, black seed, fenugreek and green tea for treatment of gastric acidity and gastric infection (84). Laurindo *et al.* (85) reported that phytochemicals can interfere with different inflammation targets such as NF- κ B pathways, and the most important chemical components for treatment of inflammatory bowel diseases are kaempferol, resveratrol, curcumin, astragalin, pinocembrin, sesamol, botulin, berberine, thymol, fisetin, gingerol, lycopene, galangin, euphol, and nobiletin. Medicinal plants which target Nrf2 pathways during treatment of inflammatory bowel diseases and related colorectal cancer are *Cynara cardunculus* L., *Panax ginseng*, *Ficus pandurata*, *Moringa oleifera*, *Ribes nigrum*, *Vaccinium myrtillus*, *Vaccinium myrtillus*, *Vitis vinifera*, *Forsythia suspensa*, and *Artemisia argyi* (85). It is reported that pomegranate has different chemical constituents such as anthocyanins, ellagic acid, punicalagin, citric acid, linolenic acid, and ascorbic acid which has important role in treatment of digestive diseases (86). Heartwing sorrel (*Rumex hastatulus*) contains ethyl acetate, n-Hexane, rutin, and aqueous fractions with anti-inflammatory and antioxidant activity, and anti-ulcer potential which can increase the level of protective glutathione-S-transferase, glutathione, and catalase and down-regulates oxidative stress marker (LPO) (87).

Gallstones

There is a high prevalence of gallstone disease in both developed and developing countries as a consequence of environmental, biochemical, and genetic factors (88-93). Gallstone disease is associated to low serum phytosterol levels, and hypersecretion of cholesterol in bile (94). It can stimulate the gallbladder mucosa and may cause acute/chronic cholecystitis and even gallbladder carcinoma (95,96). The symptoms of gallstone disease are nausea, abdominal pain, and fever, and the main subtypes of gallstones are categorized into three types: phosphate gallstones, pigment gallstones, and cholesterol gallstones (97). As a multifactorial disease which can be affected by both negatively and positively by diet, many samples of the protective impacts of dietary elements can prevent gallstone disease and enhance metabolic pathways (98-100). It is also found that Glutaredoxin-1 (*Glx1*), and Glrx1-related protein S-glutathionylation can have significant role in gallstone formation via the targeting of cholesterol metabolism (101). Arrout *et al.* (102) discovered that p-cymene of *Syzygium aromaticum* and *Nigella sativa* can be considered as alternative litholytic agent to prevent gallstone formation via their potential anti-inflammatory activity in the gallbladder. Medicinal herbs and plant utilized by patients for the treatment of gallstones in Morocco are onion, celery, parsley, fennel, cumin, caraway, carrot, coriander, date palm, red radish, prickly pear, fenugreek, origanum, saffron, rosemary, lavender, sage, thyme, cinnamon, marjoram, olive, barley, flax, black cumin, apple, pomegranate, green tea, sweet orange, and turmeric (102). Chang *et al.* (103) reported that vegetarian diets may lower symptomatic gallstone disease through cholesterol lowering.

Gastroesophageal Reflux Disease (GERD)

One of the most common diseases of the gastrointestinal system is GERD (104). It is commonly happened with other gastrointestinal motility disorders like functional constipation (105), and it is very common in young infants (106). It is associated a wide range non-esophageal such as angina, chest pain, and respiratory problems, as well as the extensive range of esophageal such as Barrett ,s disease, esophageal cancer, and esophageal ulcer

(107-115). It is discovered that the most important medicinal plants which are useful for treatment of GERD and its symptoms are *Fumaria officinalis*, *Artemisia asiatica*, *Atropa belladonna*, *Ceratonia siliqua*, *Ulmus rubra* bark, *Althaea officinalis* root and leaf, *Aloe vera* leaf gel, *Morus alba* leaf extract, *Curcuma longa*, and *Salvia miltiorrhiza* (116,117). The main mechanisms of medicinal plants and herbs in GERD mitigation are reducing acid production, soothing and protecting the esophagus, increasing digestive function, and alleviating inflammation (118). A famous classic traditional Chinese medicine is Wu Chu Yu Tang derived from the Shanghan Lun which consists of *Ziziphus jujube*, *Zingiber officinale*, *Panax ginseng*, and *Evodia rutaecarpa*, which has been found effective with similar effects to omeprazole for GERD treatment (119). Rose oil is common in Iranian traditional medicine which has high potential to preserve effective components, and it has notable influence as omeprazole in treatment of GERD (120). Zohalinezhad *et al.* (121) found that quince syrup significant enhanced symptomatic status and patient's condition.

Gastroenteritis

It is a very common paediatric disease and remains a noticeable cause of childhood mortality and morbidity world-wide (122,123). The global burden of gastrointestinal infections combines the effects of severity, incidence, and duration (124-126). *Polygonum Cillinerve* polysaccharide (PCP) is the main active ingredient of *Polygonum Cillinerve*, which can be used to treat transmissible gastroenteritis virus through the Caspase-dependent mitochondrial pathway following miR-181 transfection (127). One of the important traditional Chinese herbal medicines in China is *Cimicifugae* rhizoma, and its polysaccharide component has immune-enhancing activity, significantly transmissible gastroenteritis virus, replication, and inhibited the apoptosis lowering mitochondrial damage (128).

Gastritis

Chronic atrophic gastritis is known by erosion and inflammation of the gastric lining, which is a prevalent digestive disease and known as a precursor to gastric cancer (129,130). The main cause of gastritis is *Helicobacter pylori* infection (131). It is a persistent inflammation of the gastric mucosa which can induce to uncomfortable signs in patients (132,133), and the second cause of death from malignant disease worldwide is gastric cancer (134). Gastritis can cause symptoms such as digestive disturbances, discomfort, and pain affecting many people in both developing and developed countries (135). Hu *et al.* (136) reported that berberine, palmitic acid, co glycosides which are the main ingredient of *Coptis chinensis* Franch. (CCF) is known for its anti-inflammatory activities, and its potent heat-clearing, which can be used in treatment of gastritis. Song *et al.* (137) also found that the effective monomers of *Magnolia officinalis* var. *biloba*, and *Coptis chinensis* Franch. have special mechanism for treatment of chronic gastritis due to Coptisine, as their main optimal component. Luo *et al.* (138) also concluded that *Scleromitron diffusa* (Willd). R. J. Wang and *Scutellaria barbata* D. Don have shown therapeutic impacts on chronic atrophic gastritis by acting HIF-1, PI3K-Akt, JNK, and cancer-related and others. Liu *et al.* (139) also recommend *Bryophyllum pinnatum* (Crassulaceae) to treat gastritis because of their main active ingredients such as isorhamnetin, quercetin, beta-sitosterol, eicosapentaenoic acid, kaempferol, cubebinolide, and artemetin as they can target SRC, AKT1, MAPK1, MAPK8, and EGFR.

Ginger also showed anti-gastritis impacts by blocking the activation of MAPK/STAT3 signaling pathway, and it can restore gastrointestinal motility, reduced levels of inflammatory parameters, enhance gastrointestinal hormone levels, and attenuate gastric histopathological damage (140). *Geranii Herba*, is the traditional medicinal plant in northeast China and Korea with gastrointestinal inflammation activity, which can decrease the cell infiltration in gastric mucosa and ulcer ratio (141). Zhang *et al.* (142) found that Huangqi (*Radix Astragali*) can be suggested in treatment of chronic atrophic gastritis, and Boeing *et al.* (143) also showed that the leaves of *Vernonia condensata* Baker demonstrated gastroprotective impacts through gastrinergic pathways, and the inhibition of gastric secretion. Dhondrup *et al.* (144) reported that the Tibetan medicine Liuwei Muxiang Pills (LWMX pills) is a famous traditional Tibetan medicine which can maintain gastric function by suppressing oxidative stress and inflammation. Jinhong Tablet can treat chronic gastritis by decreasing the inflammatory cytokines indirectly or directly to play anti-inflammatory activity due to containing flavonoids contents (145).

Qi-Zhi-Wei-Tong granules can alleviate chronic gastritis, and it can also ameliorate the levels of inflammatory responses, and increased intestinal integrity (146). Qirui Weishu capsule as a famous traditional formula also recommended for treatment of chronic gastritis because of its effectiveness and safe effects (147). Han *et al.* (148)

concluded that Weierning tablet suppressed gastric mucosal cells apoptosis, and reversed intestinal metaplasia, and it inhibited the inflammation of gastric tissue. Zuojin Pill (ZJP) is a classic prescription composed of *Tetradium ruticarpum*, and *Coptis chinensis* which is found as an important therapeutic effect on chronic atrophic gastritis induced by MNNG, which may be closely associated to the inhibition of Akt/PI3K/TGF- β 1 signal pathway (149). Chaihu-Shugan-San, which is a famous Chinese prescription composed of *Rhizoma Cyperi*, *Rhizoma Chuanxiong*, *Fructus Aurantii*, *Rhizoma Chuanxiong*, *Fructus Aurantii*, *Radix Glycyrrhizae*, *Radix Paeoniae Alba*, *Pericarpium Citri Reticulatae*, *Radix Paeoniae Alba*, *Pericarpium Citri Reticulatae*, and *Radix Bupleuri* which has been used widely in treatment of different kinds of chronic gastritis due to some important components such as paeoniflorin, neohesperidin, naringin, merazin hydrate, liquiritin, isoliquiritigenin, hesperidin, glycyrrhizic acid, glycyrrhetic acid, ferulic acid, and albiflorin (150). The ethanol extract of roots and stem of *Alisma canaliculatum* which can be found in different parts of Taiwan, China, Japan, and Korea has anti-inflammatory potential, and it can suppress inflammatory responses in lipopolysaccharide (LPS)-stimulated macrophages (151).

Silymarin has anti-carcinogenic, antioxidant, and anti-inflammatory activity which can significantly reduce the development of gastric lesions, and inhibits ethanol-induced gastritis in rats (152). Pink knotweed (*Polygonum capitatum*) has both anti-inflammatory and antibacterial effects, and significant advantage in the treatment of *Helicobacter pylori*-associated gastritis (153). An important pungent in *Capsicum annuum* is called Capsaicin which can down-regulate proinflammatory cytokines in gastric mucosa, showed gastric protection against gastric lesion, decreased inflammatory infiltrate in gastric mucosa, and down-regulated cyclooxygenase-2 (COX-2) in gastric mucosa (154). *Taraxacum coreanum* Nakai has been traditionally applied for treating gastrointestinal diseases and gastric ulcer by decreasing inflammation and oxidative stress (155). The dried tuber of *Gastrodia elata* Blume (GEB) is called gastrodiae rhizome, which is called Tianma in Chinese language, could significantly enhance gastric acid secretion, and ameliorate the damages gastric glands via regulating purine metabolisms, and energy (156). Mendivil *et al.* (157) found that Capsaicin decreased inflammatory infiltrate in gastric mucosa, down-regulated proinflammatory cytokines in gastric mucosa, and modulated COX-2 in gastric mucosa.

It is reported that Eupatilin, which is the major ingredient of Artemisiae plants has shown antioxidant and anti-inflammatory activities, which can also increase gastroprotective impacts in gastric ulcer-induced rats (158). The pharmacological activities of traditional Chinese medicine and their major active ingredients on chronic gastritis are (1) repair gastric mucosa such as increase the expression of matrix metalloproteinase-2, reduce the activity of matrix metalloproteinase-9, and regulate TNFR-c-Src-ERK1/2-c-Fod pathway, (2) anti-apoptosis such as increase the ration of Bcl-2/Bax, and inhibit the AKT-Mdm2-p53 signal pathway, (3) anti-oxidation such as increase the content of SOD and GSH, and regulate Nrf2/HO-1 signal pathway, (4) improve gastric function such as inhibit gastric acid secretion, increase PGE2, G-17, PGR and PG I, and improve gastric motility, (5) anti-inflammation such as increase the level of anti-inflammatory cytokine, regulate MAPK, PI3K-Akt, NF- κ B signal pathways, and reduce the production of pro-inflammatory cytokines, (6) anti-*Helicobacter pylori*, such as increase the *Helicobacter pylori* clearance rate, and inhibit *Helicobacter pylori* infection (159).

Celiac diseases

Celiac disease is the most common intestinal chronic inflammatory disease (160,161), and it is an autoimmune disorder caused by the consumption of gluten which can negatively influence the small intestine (162,163). Gluten-free diet is one of the most important treatment strategies for celiac disease (164). Iacomino *et al.* (164) reported that *Triticum monococcum* amylase trypsin was appropriate for non-celiac wheat sensitivity patient diet, as its amylase-trypsin-inhibitors (ATIs) indicate a higher susceptibility to enzymatic hydrolysis and reduce INF- γ expression and IL-8 on tissue explants, and T-cells from celiac patients. Celiac disease develops in genetically susceptible individuals because of ingestion of gluten forming proteins discovered in cereals such as barley, rye, and wheat (165-167), and one of the notable treatments is a strict gluten-free diet (168,169). It is reported that a combination of a glutamine-specific endoprotease and prolyl endopeptidase from *Sphingomonas capsulata* which is known as latiglutenase showed insufficient advantages in phase II clinical trials, possibly due to its low enzyme activity in the gastric environment (170). The diagnosis of celiac disease is closely associated to the autoimmune response, and usually implies different tests like high serum levels of IgA, excluding selective IgA deficiency, and IgG anti-tTG, serum IgA, and IgG anti-deaminated gliadin peptide, HLA-DQ typing, and serum IgA and endomysial antibodies (171,172). Anemia, osteoporosis, hepatitis, and arthritis are among the extraintestinal manifestation of active celiac disease (173-178).

Diet therapy is an important alternative therapeutic approach which may focus on inhibition of intestinal permeability, enzymatic degradation of gluten, inhibition of the transglutaminase 2 enzyme, modulation of the immune response 1, induction of tolerance, and blocking antigen presentation (179). Wang *et al.* (180) showed that the peanut skin proanthocyanidins (PSPc) can decrease damage by suppressing signaling pathways of inflammation and oxidative stress, and it can also inhibit cell membrane damage and apoptosis in the Caco-2 celiac disease model cells. Lavoignat *et al.* (181) reported that wheat proteins can trigger immunogenic activities, and celiac disease because of their resistance to immunostimulatory and digestion epitopes. Ribeiro *et al.* (182) showed that L-theanine-gluten significantly decrease a T-cell-mediated immune response in celiac disease, and retained the functionality of gluten and the typical viscoelastic behavior. A prolyl endopeptidase was derived from the basidiomycete *Flammulina velutipes*, which could degrade the antigenic epitope of α -gliadin which triggers celiac disease (183). Some of the gluten-free flours are almond flour, coconut flour, rice flour, tapioca flour, chickpea flour, quinoa flour, buckwheat flour, sorghum flour, millet flour, cassava flour, soy flour, potato flour, teff flour, green banana flour, chestnut flour, sesame flour, and chia flour (184).

Peptic ulcer disease

Peptic ulcer is a chronic disease which caused by the infection of *Helicobacter pylori* (185-187), and this recurrent disease, massively influence on the mortality and morbidity of many people (188-189). Peptic ulcer can be treated by various classical clinical experience formulas, Chinese medicine decoctions, and chemical constituents made from plant sources and naturally occurring components (190,191). Peptic ulcer is an acid-induced lesion of the digestive tract which is commonly located in proximal duodenum or stomach, which is characterized by denuded mucosa with the defect extending into the submucosa or muscularis propria (192). The anti-peptic ulcer effects are gastric secretion, cytoprotective activity, angiogenesis promotion, amino acid metabolism, antioxidant activity, anti-inflammatory activity, and gastrointestinal motor activity (193-195). It is because of diminished mucosal defenses, and imbalances in gastric juice pH (196,197). Kuna *et al.* (198) found that alternative therapy for peptic ulcer treatment can be recommended and suggested medicinal plants are Korean red ginseng, *Allium sativum*, *Cistus laurifolius*, *Zingiber zerumbet*, *Zingiber officinalis*, *Camellia sinensis*, *Artemisia asiatica*, and *Curcuma longa*.

The main components of NPK500 capsules which is a *Cassia sieberiana* DC.-based herbal analgesic medicine are betulinic acid, kaempferol, and physcion which possess anti-inflammatory and analgesic activities with significant capability of relief inflammation and acute pain with causing hepatic, renal or gastrointestinal injuries (199). Kumadoh *et al.* (200) observed the positive effects of Brimstone tree (*Morinda lucida* L.), *Cnestis ferruginea* Vahl ex D.C., *Vernonia amygdalina* Del., *Momordica charantia* Linn., *Lantana camara* Linn., *Trema orientalis* Linn., *Psidium guajava* Linn., *Persea americana* Mill., *Spondia mombin* Linn., *Citrus aurantifolia* Linn., *Bidens Pilosa* L., *Maytenus senegalensis* Lam., *Carapa procera* D.C., *Trichilia monadelpha*, and *Cassia sieberiana* D.C. in the management of peptic-ulcer disease. Huang *et al.* (201) reported that two Chinese herbal formula namely Hai-Piao-Xiao and Ban-Xia-Xie-Xin-Tang could be suggested for treatment of peptic ulcer diseases in countries such as China and Taiwan. Photogedunin and gedunin of *Xylocarpus granatum* has shown anti-ulcerogenic activity which may be because of its anti-secretory activity with positive effects against peptic ulcer disease (202). One of the famous Indian traditional medicinal plants is *Anogeissus pendula* Edgew (Combretaceae) with high antioxidant activity, and positive effects in treatment of peptic ulcer disease (203). Jesus *et al.* (204) recommended *Hyptis suaveolens* (L.) Poit (Lamiaceae) for application as a gastro protective agent in central region of Brazil. *Piper umbellatum* L. (Piperaceae) is a shrub discovered in Atlantic Forest region, Savannah, and the Amazon of Brazil which is recommended for the treatment of gastric disorders due to its high rutin, quercetin, and flavonoids (205).

Another medicinal plant from Brazil is *Uncaria tomentosa* which can prevent piroxicam- and ethanol-induced ulceration as it has high anti-inflammatory, and antioxidant activities (206). B-Myrcene and some other major components of *Citrus aurantium* could enhance in mucosal malondialdehyde level with significant anti-ulcer activity (207). Memariani *et al.* (208) reported that the beneficial effects of *Pistacia atlantica* oleoresin which contains α -pinene, as complementary medicine and dietary supplement in folk traditional medicine for the treatment of PU. Different components of *Tagetes erecta* L. such as oil, disc florets, and ray can decrease the risk of ulcers, and it has also potential to cure acid pepsin disorders (209). Its extract can also accelerate the gastric healing process, decrease gastric inflammation, and reduce the ulcer recurrence (210). Khan *et al.* (211) showed that the extract of *Mangifera indica* possess significant impact on amelioration of gastric ulcer by re-inforcing the gastric mucosa,

preventing of indomethacin-induced muco-oxidative stress, and decreasing the secretion of gastric acid. Some of the most important medicinal plants in Nigerian traditional medicines for treatment of peptic ulcer are leaves of *Synclisia scabrida*, the whole plant of *Diodia sarmentosa*, leaves of *Ficus exasperate*, and leaves of *Cassia nigricans*, which reduce both the acidity and volume of gastric secretion, enhanced the pH, and delayed intestinal transit (212).

Bittersweet orange (*Citrus x aurantium* var. bigaradia) which is an old hybrid between *Camellia reticulata* and *Citrus maxima* from China with more than 4000 components, especially flavonoids, moreover, it has also contained neohesperidin and hesperidin which can notably aggravate gastric damage by histopathological changes, and enhances ulcer index (213). Stemmoside C of *Solenostemma argel* which is widely distributed in Asia and Africa. It can be used against gastric ulcer progression and tumorigenicity (214). It has been showed that luteolin and kaempferol extract of *Kaempferia galanga* L. galanga have protective impacts against ethanol-induced gastric ulcer (215). Figueiredo *et al.* (216) also reported that the chemical components of *Fridericia chica* L.G. Lohmann (Bignoniaceae) can be used to cure gastrointestinal disorders such as peptic ulcer. Liuwei Anxiao San which is one Mongolian medicinal prescription can be used against gastric mucosal injury in gastrointestinal disorders by suppressing the JAK2/STAT2 pathway (217). In Malay community, *Dicranopteris linearis* (DL) known as resam with high antimicrobial and antioxidant activity, and contains saponins, terpenoids, and flavonoids which is recommended for application for treatment of peptic ulcer disease (185). De Jesus *et al.* (218) tannins are poly-phenols presents in beverages, foods, and plants with potential ability for treatment of peptic ulcers. *Ranunculus millefoliantis* extract revealed high gastroprotective impacts that could be because of decrease malondialdehyde quantity, rise superoxide dismutase and catalase, capability upsurge pH then mucus discharge, down-regulating Bcl-2-associated X protein protein, up-regulating heat shock protein 70, and moderate provocative cytokines (219).

Ruscogenin (C₂₇H₄₂O₄) is a natural steroid sapogenin product derived from *Tribulus terrestris* with antioxidant and anti-inflammatory activity and suitable for treatment of peptic ulcer (220). Wu *et al.* (221) found that the burdock essence can help abolish *Helicobacter pylori* infection, and increase the therapeutic impacts of conventional medication on gastric mucosal repair in gastrointestinal ulcer patients. *Adinandra nitida*, which is known as Shibi tea, is rich in flavonoid and considered as an important medicinal plant in South China, moreover, Shibi tea water extract (STE), and its major flavonoid camellianin A (CA) can regulate the inflammatory signaling pathways protein levels of NF-κB and IκB-α (222). Mohammad *et al.* (223) noted that henna (*Lawsonia inermis* L.) is effective, low cost, safe and suitable for treatment of gastric ulcer. Orsi *et al.* (224) reported that the methanolic extract obtained from the stem bark of *Hymenaea stigonocarpa* Mart. Ex Hayne (Fabaceae) showed significant gastroprotective, anti-diarrheal, and cicatrizing effects which may be because of the presence of flavonoids and condensed tannins in the fruit and bark of the plants. Propolis is an important natural component which has been used for many years in folk medicine, and it refers to bee glue consists of leaves, plant resins from saps, barks, and buds of plants collected by bees around their vicinity, and its application can enhance the amounts of catalase, superoxide dismutase, glutathione, nitrite, while malondialdehyde contents may decrease which proves their important for management of peptic ulcers diseases (225). *Celosia trigyna* belongs to the plant family Amaranthaceae which is found in Nigeria, Southern Arabia, South Africa, and Congo with anti-ulcerogenic effects, and its hexane extracts, and secondary metabolites such as sterols, pheophytin a, and lutein exhibited high effectiveness in treatment of gastric ulcer (226). Algandaby (227) concluded that *Teucrium polium* extract muco-adhesive formula can be considered as an important therapy for chronic oral ulcers, as its application could significantly decrease TNF-α, MDA, COX-2, and IL-6, and increase catalase and superoxide dismutase activities. The root extracts of *Prangos pabularia* (REPP) application can meaningfully modulated oxidative stress, increase MDA, PGE2, CAT, and SOD, and inflammatory cytokines levels which introduces it as an important remedy for gastric ulcer disorder (228). Hasna *et al.* (229) suggested that fenugreek extract combined with *Bifidobacterium breve* can be an alternative for treatment of diseases caused by *Helicobacter pylori* infection according to Algerian medicinal plant sciences. *Justicia pectoralis* Jacq. which is an important medicinal plant in several Latin American countries such as Brazil has antioxidant and anti-inflammatory activities, and its leaves methanol extract can be used for gastric disorders (230).

In African countries, especially in South Africa, medicinal plants such as *Garcinia kola*, *Sclerocarya birrea*, *Combretum molle*, *Strychnos species*, and *Alepidea amatymbica* have been used for several years for treatment of peptic ulcer and gastritis disease (231). Zhang *et al.* (232) observed that the *Dendrobium officinale* polysaccharide

(DOP) ameliorated gastric ulcer through considerable improve in expression of SOD in tissues, and SOD, GSH, and GSH-PX in serum. In Ethiopia, medicinal plants which have been used for treatment of gastric ulcer are *Urtica simensis*, *Croton macrostachyus*, *Achyranthes aspera*, *Afrocarpus falcatus*, *Aloe gilbertii*, *Aloe macrocarpa*, *Aloe pubescens*, *Asparagus africanus*, *Asystasia gangetica*, *Brassica carinata*, *Brassica oleraceae*, *Calpurnia aurea*, *Carica papaya*, *Carissa edulis*, *Casimiroa edulis*, *Catha edulis*, *Caylusea abyssinica*, *Centaurium pulchellum*, *Cicer arietinum*, *Citrus aurantifolia*, *Citrus sinensis*, *Commiphora hodai*, *Commiphora kua*, *Cordia Africana*, *Croton macrostachyus*, *Cucurbita pepo*, *Dodonaea Angustifolia*, *Enset ventricosum*, *Erythrina brucei*, *Euphorbia abyssinica*, *Flacourtia indica*, *Galium simense*, *Girardinia bulbosa*, *Hibiscus cannabinus*, *Justicia schimperiana*, *Kalanchoe petitiiana*, *Leonotis ocymifolia*, *Lepidium sativum*, *Lippia adoensis*, *Momordica foetida*, *Nigella sativa*, *Ocimum lamiifolium*, *Oncoba spinosa*, *Physalis peruviana*, *Plantago lanceolata*, *Psidium guajava*, *Rubus apetalus*, *Rumex nepalensis*, *Rumex nervosis*, *Salvia merjamie*, *Cordia Africana*, *Rumex nepalensis*, *Osyris quadripartite*, and *Plantago lanceolata* (233). Medicinal plants and herbs in Indian traditional medicinal science for treatment of peptic ulcer are *Jasminum grandiflorum*, *Anogeissus latifolia*, *Alchornea castaneaeifolia*, *Uleria salicifolia*, *Solanum nigrum*, *Ocimum sanctum*, *Scoparia dulcis*, *Byrsonima crassa*, *Asparagus racemosus*, *Centaurea solstitialis*, *Anacardium occidentale*, *Calophyllum brasiliense*, *Larrea divaricata*, *Hemidesmus indicus*, and *Spartium junceum* (234).

Colorectal polyp

A colorectal polyp is a polyp happening on the lining of the rectum or colon (235), and untreated colorectal polyps can develop into colorectal cancer (236,237). It is usually categorized by its cause and the behavior (238), and they maybe malignant, pre-malignant, or benign (239). Colorectal polyps refer to tissue on the surface of the large intestinal mucosa which is shaped by abnormal hyperplasia of the epithelium (240-242). Polyps are categorized into two major groups, namely adenomatous and hyperplastic polyps (243,244). Various natural components can suppress or induce apoptosis proliferation of tumor cells (245,246). Flavones which have shown anti-colorectal cancer activity are baicalein, apigenin, and digitoflavone (247). Herbal plants and medicinal plant extracts can prevent the growth of colorectal cancer by activating several signaling pathways, cell cycle arrest at different stages, and inducing autophagy or apoptosis (248-250).

Traditional medicinal herbs and plant which have been reported useful for treatment of colorectal polyp are *Curcuma longa*, *Panax quinquefolius*, *Olea europaea* var. *sylvestris*, *Garcinia mangostana*, *Emilia sonchifolia*, *Rabdosia rubescens*, and *Hylomecon vernalis* (251). Flavonoids are considered as the secondary metabolites which have shown high anti-colorectal cancer effects (252-255). Phytochemicals can modulate the production of cytotoxic components such as ROS, and NO by macrophages, moreover, they can polarize macrophages towards the M1 phenotype, inhibit M2 polarization, and increase anti-tumor immune responses (256). High intake of hydroxycinnamic acids, total phenolic acids, and flavonols could also decrease the risk of colorectal cancer (257). The main bioactive ingredient of *Cuminum cyminum* is cuminaldehyde which is also known as 4-isopropylbenzaldehyde which has beneficial effects against human colorectal and cervical carcinoma (258,259). Curcumin has been derived from dried roots of the plant *Curcuma longa* which can be used for the treatment of colorectal cancer (260). It is also proved that apigenin is capable to decrease polyp numbers as well as increasing protein 53 activation via phosphorylation in animal models (261). Cao *et al.* (262) also found the importance of application of berberine in treatment of colorectal disease through multiple targets and channels. Ansari *et al.* (263) reported the inhibitory effects of naringin on colorectal disease via different signaling pathways. Cai *et al.* (264) reported that Xianlian Jiedu Decoction can be suggested for treatment of colorectal disease by reducing metabolic disorders, intestinal microbial flora, and alleviating the level of inflammation. Fernandez *et al.* (265) reported that some important flavonoids such as luteolin and apigenin can be for treatment of colorectal cancer cell lines due to its antitumor activity. Cassotta *et al.* (266) also found that ω -3 polyunsaturated fatty acids, vegetables, and fiber can prevent inflammatory bowel disease related colorectal cancer.

Drimys calcarata bulb extract showed high tannin, total phenolic and flavonoid content as well as significant antioxidant activity, and the methanol extract induced mitochondrial-mediated apoptosis in colorectal HT-29 and Caco-2 cells with beneficial effects for treatment of colorectal cancer (267). Ling *et al.* (268) also showed that berberine can influence macrophages polarization and infiltration, and inhibited tumorigenesis in colorectal cancer. In South America, cocoa powder is found important to prevent colorectal cancer through modulating oxidative

stress (269). It is reported that chemical components of the most important medicinal plants in traditional Chinese medicine like *Smilax china*, *Patrinia scabiosaefolia*, and *Sanguisorba officinalis* can be recommended for treating colorectal cancer as HSD11B2 is the main target protein of them in treating colorectal cancer (270). Nascimento *et al.* (271) also reported that jaboticaba peel is an important source of phenolic acids and cyanidin-3-*O*-glucoside with high antioxidant activity which could mitigate colorectal cancer probably by avoiding triggering inflammation. Khan *et al.* (272) concluded that the alcoholic leaf extract of *Annona reticulata* has anti-proliferative and antioxidant activity, which can improve prevention of colorectal cancer development and dysplasia caused by 1,2-dimethylhydrazine. Nomura *et al.* (273) reported that the intake of fruits and vegetables was inversely connected to colorectal cancer risk among men. Nirmala and Ramanathan (274) reported that kaempferol could be notably applied as chemo-preventive agent in colorectal cancer. *Frangula alnus* ethyl-acetate extract (FA), and its dominant component emodin (E) demonstrated high cytotoxic activity against carcinoma cells, modulated cell cycle, and induced cell cycle (275). Polyhydroxybutyrate is a non-toxic polyhydroxyalkanoate polymer produced by different microorganisms which has been reported to increase cardiac performance, enhance cognitive and brain function, and significantly regulate diabetes and appetite suppression for treatment of colorectal cancer (276). Wang *et al.* (277) reported that naringenin is an important component for inhibiting colorectal cancer proliferation, probably via the AMP-activated protein kinase (AMPK) pathway, to induce apoptosis in colorectal cancer and regulate mitochondrial function. Medicinal plants with anti-colorectal cancer bioactive ingredients with high potential in colorectal cancer management are *Musa balbisiana*, *Musa acuminata*, *Punica granatum* L., *Glycine max*, *cruciferous vegetables*, *Hibiscus tosa-sinesis*, and *Hibiscus sabdariffa* L. (278).

Bloating

Among disorders of gut-brain interaction, bloating is a common symptom in the general public (279-281). Abdominal distension and bloating are secondary to an organic disease (282), which can influence social interactions, daily life, and can be triggered by dietary habits (283,284). Bloating is the result of gas release in presence of the notable proper amount of liquid phase with a suitable viscosity (285). Abdominal bloating and distension can develop for different reasons, including a previous infection, food intolerance, delayed intestinal transit, disordered visceral sensation, and an abnormal viscero-somatic reflux (286,287). Application of *Bacillus coagulans* GBI-30, 6086 probiotic can be suggested as it is effective and safe for the relief of bloating and abdominal pain for the patients with symptoms of irritable bowel syndrome (IBS) (288). KAASER is one of the most important Persian herbal formulations comprised of *Piper nigrum* L. berry, *Zingiber officinale* Roscoe. rhizome, and *Trachyspermum ammi* (L.) Sprague seed which is effective to treat patients suffering from bloating and flatulence (289).

It is also reported that usage of vitamin D, psyllium, and probiotic supplementation can be useful in treatment of bowel syndrome (290). It is also discovered that a novel therapeutic of natural origin which contains a mixture of naturally occurring polysaccharides, tannins derived from grape seed extract, pea proteins and β -galactosidase can be used in treatment of distension, bloating, and abdominal pain in patients with lactose intolerance (291). Patch *et al.* (292) reported that application of *Bacillus subtilis* (BG01-4TM) can increase specific symptoms of constipation and associated gastrointestinal dysfunction. Bertin *et al.* (293) also emphasized on the beneficial effectiveness of fermented monosaccharide, disaccharide, oligosaccharide, and polyol diet for patients with irritable bowel syndrome. Extracts of *Boswellia serrata* and *Curcuma longa* have antimicrobial and anti-inflammatory effects and their application can significantly reduce irritable bowel syndrome (294).

Pancreatitis

Acute pancreatitis is an inflammatory disease of the exocrine pancreas (295,296) with significant mortality and morbidity (297-299), which is characterized by substantial necrosis, vacuolization, and interstitial edema (300-302). It was a common acute abdominal disease known by inflammation and pancreatic acinar cell death (303,304). Ramakrishnan *et al.* (305) introduced curcumin, rhein, green tea catechin derivatives, resveratrol, emodin, ellagic acid, embelin, eruberin A, and metformin as important anti-fibrotic agents for pancreatic cancer and chronic pancreatitis. Chemical structures of the phytochemicals which show protective effects against acute pancreatitis are withaferin A, Visnagin, Tetramethylpyrazine, Tanshinone IIA, Saikosaponin a, rutin, resveratrol, quercetin, picroside II, oxymatrine, mogroside IIE, luteolin, curcumin, ellagic acid, dihydrokaempferol, emodin, chlorogenic acid, borneol, berberine, carvacrol, calycosin, baicalin, baicalein, artesunate, and acetyl-alpha-boswellic acid (306).

As an important natural flavonoid component which can be found in different vegetables, fruits, and plants, myricetin can alleviate the severity of acute pancreatitis via inhibition of nuclear factor- κ B, and cathepsin B activity, and myricetin can be applied as a regulator of Ca^{2+} /CaM-dependent protein kinase kinase 2 (CaMKK2), and calcineurin signaling (307).

It is reported that tectoridin which is an isoflavone, and a type of flavonoid can ameliorate pancreatic injury and inflammation in severe acute pancreatitis mice, as it can target ERK2 to inhibit ERK MAPK signaling pathway (308). Tropolone is an organic component with the chemical formula $\text{C}_7\text{H}_5(\text{OH})\text{O}$, and it is reported that tropolone derivative hinokitiol as an important phytochemical with anti-inflammatory activities can reduce the expression of COX-2, and NF- κ B, and reduce the expression of proinflammatory cytokines (309). Chaikin chengqi decoction can reduce acute pancreatitis by targeting gasdermin D-mediated pyroptosis, and then is the main active ingredient of it in suppressing pyroptosis (310-312). Wen *et al.* (313) reported that the effectivity of Chaikin chengqi decoction is mainly because of its ingredients such as chrysin, rhein, baicalin, emodin, and ingredients. Han *et al.* (314) reported that Qing Xia Jie Yi Formula (QXJYF) is an important herbal formula from traditional Chinese medicine which can alleviate acute pancreatitis via inhibition of M1 macrophage polarization by suppression glycolysis. Baicalin is a flavonoid component derived from the root of *Scutellaria baicalensis* Georgi, which is known as *Huangqin*, with anti-inflammatory activity, and it can suppresses oligomerization of phosphorylated mixed lineage kinase domain-like protein (MLKL) by decreasing ROS, and it can also mitigate the severity of cerulein-induced acute pancreatitis in mice (315). Abozaid *et al.* (316) reported that Cinnamic acid nanoparticles (CA-NPs) can be applied as a modulator for the redox signaling pathway of pancreatitis with both anti-inflammatory and antioxidant activity. Nimbolide as the main limonoid of the neem tree, is extensively applied in traditional medicinal treatment in South of Asia with antitumor, anti-inflammatory, and antioxidant activities, and it can protect against cerulein-induced acute pancreatitis, reduce pancreatic inflammation through regulating NF- κ B/SIRT1 pathway, and suppress pancreatic apoptosis by modulating cleaved Bcl2, Bax, and caspase 3 expression (317).

Hesperidin is a flavanone glycoside obtained from citrus fruits, and it can provoke deficit in glucose control and insulin secretion, and prevent pancreatic inflammation through increasing antioxidant mechanism (318). Fan *et al.* (319) reported that Wedelolactone can attenuate pancreatitis and some other lung injury in Taurocholate or Cerulein-induced murine models with ability of suppression of ferroptosis and pyroptosis through the up-regulation of glutathione peroxidase-4 (GPX4), as it is a polyphenol obtained from *Eclipta alba*, and *Wedelia chinensis*. Dachengqi decoction (DCQD), is a famous traditional Chinese medicinal formula which consists of *Magnoliae officinalis cortex*, *Aurantii fructus immaturus*, *Natrii sulfas*, and *Rhei radix ex rhizome*, and effective for treating inflammation in different disease conditions, and abdominal pain (320). In another experiment, it is showed that Matrine, which is a natural alkaloid obtained from stems and roots of *Sophora flavescens*, can alleviate ferroptosis and oxidative stress in severe acute pancreatitis (321). One of the most important polyphenols with anti-inflammatory activity is Chlorogenic acid which can significantly influence L-arginine-induced pancreatitis and pancreatitis-associated lung injury (322).

Dihydrokaempferol is one type of flavonoids which can be obtained from *Bauhinia championii* (Benth) which can regulate the protein Kelch-like ECH-associated protein (Keap1)/the nuclear factor erythroid 2-related factor 2 (Nrf2) pathway and regulating oxidative stress injury in severe acute pancreatitis (323). Guo Qing Yi Tang is composed of 13 traditional Chinese medicines such as senna leaf, sodium sulfate, raw rhubarb, *Magnolia officinalis*, hawthorn, and *Bupleurum*, which can reduce the level of inflammatory factors, and enhance intestinal mucosal barrier function for treatment of acute pancreatitis (324). Hog plum (*Ximenia americana*) contains heteropolysaccharide with antinociceptive impacts, which can decrease pancreatic enzymes as well as inflammatory parameters of acute pancreatitis (325). *Morus alba* root bark extract has high anti-inflammatory activity due to the presence of Cudraflavone B which can be suggested for treatment of pancreatitis (326). Lima *et al.* (327) reported that 1,8-cineole can attenuate cerulein-induced acute pancreatitis through an anti-inflammatory process, and by combating oxidative stress. Zhang *et al.* (328) in his experiment showed that the extract of *Ligustrum vulgare* L. significantly decreased the fibrotic changes and the inflammatory activity which proves the potent pancreato-protective impacts of it. *Nigella sativa* Linn. is a medicinal plant which is rich in chemical components with anti-inflammatory activity possibly by managing and regulating the expression of ASC protein of the Nod-like Receptor Pyrin domain containing 3 (NLRP3) inflammasome in pancreas that may reduce the activation of caspase-1 (329).

It is also reported that the usage of apigenin to humans could significantly ameliorate the damages to the pancreas (330). Seo *et al.* (331) reported that quercetin 3-*O*-xyloside has significant anti-pancreatitis activity via attenuating intracellular ROS production and ER stress response, and increasing apoptotic cell death. One of the most important phenolic components of long pepper (*Piper longum*), and black pepper (*Piper nigrum*) fruits which can decrease the digestive enzymes and pro-inflammatory cytokines, and it can significantly inhibit the activation of mitogen activated protein kinases that may show its effectiveness as a protective effect against acute pancreatitis (332). Fisetin is a flavonoid contained in different foods such as fruits, teas, wines, and juices which can inhibit the activation of c-Jun NH2-terminale kinase (JNK) and nuclear factor (NF)- κ B *in vitro* and *in vivo* with high anti-inflammatory impacts on acute pancreatitis (333).

Chinese herbal ointment Liu-He-Dan which is composed of *Angelica dahurica*, *Phellodendron chinense* Schneid., *Rheum officinale* Baill., *Armeniaca mume* Siebold, flour, honey, and *Mentha haplocalyx* Briq., could be applied to relieve the disease of pancreatitis (334). Anchi *et al.* (335) observed that curcumin which is a polyphenolic derivative from turmeric (*Curcuma longa*) could significantly reduce lipase levels, serum amylase, nitrosative and oxidative stress, and appropriate for treatment of pancreatitis through sustained release microparticle approaches. Choi *et al.* (336) found that berberine inhibited cerulein-induced acute pancreatitis, exerted anti-inflammatory effects, and inhibited inflammatory mediators production in acute pancreatitis. Genistein which is a polyphenolic isoflavone and belongs to the flavonoid in vegetables such as fava beans, and soy beans attenuated cerulein-or taurocholate-induced acute pancreatitis *in vivo*, and promoted apoptosis of pancreatic acinar cells to protect from acute pancreatitis (337). There is a compound and plant steroid in *Commiphora myrrha*, which is known as Guggulsterone, protected lung injury and the cerulein-induced pancreatic, inhibited the activation of ERK1/2 and JNK in cerulein-induced acute pancreatitis, and inhibited cerulein-induced elevation macrophage and neutrophil infiltration (338). Bansod *et al.* (339) concluded that berberine activated AMPK and downregulated TGF- β /Smad signaling pathway, and attenuated cerulein-induced chronic pancreatitis and associated fibrosis. Kumar and Tamizhselvi (340) also indicated that garlic-derived organosulfur component diallyl disulfide (DADS) alone or combination inhibited NF- κ B translocation, and I κ B degradation. It is reported that flavonoid C1 from *Coreopsis tinctoria* Nutt. has potential role in treatment of acute pancreatitis through Nrf-2/ARE-mediated antioxidant pathways and due to its antioxidant effects (341).

Moreover, aqueous extract of black tea (*Camellia sinensis*) prevents pancreatitis caused by chronic ethanol (EtOH)+cholecystokinin (CCK) toxicity possibly by increasing antiapoptotic, anti-inflammatory, and antioxidant activity in rats (342). Bashir *et al.* (343) reported that grape seed proanthocyanidins has an effective function as a protector in the oxidative stress-mediated pancreatic dysfunction in rats. One of the most important traditional medicinal plants in Brazil is *Caesalpinia pyramidalis* Tul. from Fabaceae family, and its ethanol inner bark extract could decrease hyperalgesia, lipoperoxidation, and inflammation in common bile duct obstruction (CBDO)-induced acute pancreatitis (344). Visnagin is an antioxidant furanocoumarin component of the plant *Ammi visnaga*, which can decrease inflammatory cytokines in intestine and lungs, and regulated the expression of markers of multiple organ dysfunction syndrome that may prove its effectiveness to prevent cerulein induced acute pancreatitis (345).

Functional dyspepsia

It is commonly known as chronic indigestion which is an upper gastrointestinal disorder (346), and associated complaints may include vomiting, nausea, belching, bloating, etc. (347). Functional dyspepsia is a complicated symptoms referable to the gastroduodenal region of the gastrointestinal tract which consists of burning or pain, early satiety, or postprandial fullness (348-350). Dyspepsia is known as episodic or persistent abdominal pain or discomfort of the upper gastrointestinal tract (351), and it has been linked with a deficiency or dysfunction of digestive enzymes (352). It is an extremely common disorder of gut-brain interaction which can influence the quality of life without an associated increase in mortality (353,354). It most usually presents with bothersome signs after eating (355,356). The positive effects of traditional Chinese medicine and compounds in the treatment of functional dyspepsia have been reported by several researches (357-359).

Mechanisms of traditional Chinese medicine components in treating functional dyspepsia are reduce visceral hypersensitivity for Si-Ni-San, modified Liu-Jun-Zi decoction; relieve anxiety and stress for Zhizhu Kuanzhong capsule, Qi-Zhi-Wei-Tong granules, and Xiangsha Liujunzi granules; regulating brain-gut peptides for Chaihu

Shugan San, Banha-Sasim-Tang, and Xiangsha Liujunzi granules, reduce low-grade duodenal inflammation for Si-Ni-San, promote gastric motility for Pixu I recipe Shugan Hewei decoction, and regulating the gastrointestinal microbiota for Shen-Ling-Bai-Zhu-San, Jiao Sanxian, and Xiaoyaosan (360,361). Saikosaponin D is the basic active monomer of saikosaponin which can improve intestinal cells of Cajal structure and morphology, that may inhibit excessive autophagy, and increased a gastrointestinal motility disorder and functional dyspepsia via regulating substance P and ghrelin levels (362). Liu *et al.* (363) concluded that the application of laminarin in different traditional medicines can regulate somatic and visceral hypersensitivity. Simo Decoction is a famous traditional Chinese medicine which composed of *Radix linderae*, *Semen arecae*, *Radix Aucklandiae*, and *Fructus aurantia*, which can improve the production of short-chain fatty acids for alleviation of functional dyspepsia (364). Xiaopi Hewei Capsule which is a classic formula comes from the Chinese Miao minority, which can significantly reduce the symptoms of functional dyspepsia (365). Xiangsha-Liujunzi-Tang is a classical formula for treatment of some digestive disease systems which can reduce the inflammation, and increase the integrity of duodenal mucosa (366). *Cynanchum auriculatum* Royle ex Wight (CA) is a perennial herbaceous plant in Hunan, Sichuan, and Guizhou of China which can be used to treatment of functional dyspepsia with significant effect on ATP-binding cassette transporter inhibitors (367).

Another important traditional Chinese medicine is Wei-Tong-Xin which is the combination of *Aucklandia lappa* Decne, *Rheum palmatum* L., *Pharbitis nil* (L.), *Gleditsia sinensis* Lam. and *Glycyrrhiza uralensis* Fisch., and its application can inactivate the toll-like receptor 4 (TLR4)/MyD88 signaling pathway and inhibit the occurrence of gastric antral inflammation (368). Moghtaderi *et al.* (369) reported a significant connection between functional dyspepsia and stomach dystemperaments after application traditional Persian medicine. Shen-Ling-Bai-Zhu-San which includes *Atractylodes macrocephala*, *Poria cocos*, and *Panax ginseng* could ameliorate dyspepsia, and improve the dysregulated composition and function of the gut microbial community. Ho *et al.* (370) also reported that Modified Ban Xia Xie Xin Decoction, and Xiao Yao Pill can be used instead of chemical drugs with effective impacts, and as alternative medicine for treatment of functional dyspepsia. Gan *et al.* (371) also reported that Ban Xia Xie Xin was more effectual in treatment of functional dyspepsia than Western Medicine. Mohtashami *et al.* (372) also showed that the application of honey-based formulation of black cumin (*Nigella sativa*) seed oil can cause meaningful symptomatic improvement of patients with functional dyspepsia. Pennyroyal (*Mentha pulegium*) extract could eradicate *Helicobacter pylori*, enhance dyspeptic symptoms, and the quality of life in functional dyspepsia patients (373). The extract of *Poria cocos* could increase the effects on functional dyspepsia through multiple pathways (374).

The herbal medicine STW5 which is a composed of drug extracts of milk thistle, Angelica root, chamomile flower, liquorice root, peppermint leaf, lemon balm leaf, and a fresh plant extract of bitter candytuft (*Iberis amara*) and ethanol content of 31% is recommended for treatment of irritable bowel syndrome and functional dyspepsia (375). ZhiShiXiaoPi Tang (ZSXPT) which is a traditional Chinese medicine includes 10 Chinese traditional medicine plants including *Magnolia officinalis*, immature bitter orange, *Pinellia ternate*, *coptis*, malt, *Rhizome zingiberis*, *Poria cocos*, *Rhizome atractylodis macrocephalae*, liquorice, and *Codonopsis pilosula* can be used against autophagy induced by corticosterone and functional dyspepsia basically by blocking the mTOR pathway in Cort-induced PC12 cells in rats with functional dyspepsia (376). Sufoof-i-Tabkheer is a powder dosage from containing *Elettaria cardamomum*, *Bambusa arundinaceae*, *Coriandrum sativum*, and *Foeniculum vulgare* can be used in treatment of dyspepsia in traditional Iranian medicine (377). Wang *et al.* (378) also reported that Chaihu-Shugan-San as an important and classic traditional Chinese medicine formula is significantly effective in treatment of functional dyspepsia, increases gastric emptying rate, and relieves depression and anxiety in patients. Jin *et al.* (379) reported that silver nanoparticles stabilized by chitosan-polyvinyl alcohol polymers mediated by *Pistacia* extract can be used for treatment of gastric cancer, and functional dyspepsia. Pingwei capsules includes *Rhizoma Atractylodis Lanceae*, *Cortex Magnoliae Officinalis*, *Radix Aucklandiae*, *Fructus Aurantii Submaturus*, *Radix Bupleuri Chinensis*, *Radix Paeoniae Alba*, *Rhizoma Bletillae Striatae*, *Endoconcha Sepiellae*, *Rhizoma Sparganii*, *Bulbus Fritillariae Thunbergii*, *Rhizoma Coptidis*, *Herba Taraxaci Mongolici*, *Rhizoma Corydalis Yanhusuo*, and *Rhizoma Curcumae Phaeocaulis* which can improve gastrointestinal motility in rats with functional dyspepsia (380). Indian Gooseberry and Amal (*Phyllanthus emblica* L.) is a traditional medicine in subtropical and tropical regions like Malaysia, China, India, Pakistan, etc., which is rich in vitamin C, amino acids, flavonoids, fatty acids, sterols, terpenes, tannins, phenolic acids, and alkaloids which can regulate the microbiome structure, and restore homeostasis by stimulating important bacterial abundance, and reduce the negative effects of bacterial abundance (381). Choi *et al.* (382) revealed that *Foeniculi fructus* which is an important traditional medicine in Europe and

China can provide a potential means to treat functional dyspepsia due to its key components such as β -sitosterol, oleic acid, and 12 functional dyspepsia-related genes. Yukgunja-tang (Liujunzi-tang in Chinese, Rikkunshito in Japanese) is an important traditional herbal medicine in East of Asia which includes *Zizyphi Fructus*, *Zingiberis Rhizoma Recens*, *Glycyrrhizae Radix et Rhizoma*, *Poria Sclerotium*, *Atractylodis Rhizoma Alba*, *Ginseng Radix*, *Citri Unshius Pericarpium*, and *Pinelliae Tuber* which also showed high safety, the effectiveness in the treatment of functional dyspepsia (383).

It is reported that Xiang Sha Liu Junzi decoction can promote the gastrointestinal motility disorder in functional dyspepsia with spleen deficiency syndrome which is in connection with reconstruction of the mitochondrial quality control system by restraining PINK1/Parkin-mediated mitophagy and division (384). One of the most important medicinal plants in China is *Amomum tsaoko* Crevost et Lemarie which is a perennial herbaceous plant belonging to the *Zingiberaceae* family which has been used to treat abdominal discomfort, postoperative abdominal distension, and very useful in treatment of functional dyspepsia (385). Therapeutic effects of traditional Chinese medicine on functional dyspepsia include treatment for smoothing liver and regulating stomach, treatment for strengthening spleen and promoting qi stagnation, treatment for acrid dispersing and bitter descending, treatment for eliminating heat and removing dampness, and treatment for warming stomach and invigorating spleen (386). Representative prescriptions in five syndrome types are Chaihu Shugan Powder for liver-stomach disharmony; Simo Decoction, Xiangsha Liujunzi Decoction, and Zhizhu Pills for spleen deficiency with qi stagnation; Banxia Xiexin Decoction for syndrome of intermingled heat and cold; Lianpo Drink and Sanren Decoction for dampness-heat of spleen and stomach; and Huangqi Jianzhong Decoction for deficiency-cold of spleen and stomach (387). It is also reported that application of *Ocimum basilicum* can be appropriate to relieve the symptoms of functional dyspepsia particularly in young and female patients with dysmotility (388).

Malabsorption

Malabsorption can cause a large variety of gastrointestinal signs such as abdominal pain, steatorrhea, constipation, diarrhea, and malnutrition (389,390). As a common clinical condition, it can influence different races around the world because of its structural and functional abnormalities of the intestinal mucosa which make it unable to absorb the nutrients in the diet (391,392). It can be caused by different diseases of the small intestine, biliary tract, diseases of the liver, pancreas, and stomach (393-395).

Bowel obstruction

Bowel obstruction is known as the main common surgical emergency following abdominal procedures (396,397). It is more common in the small bowel than in the large bowel (398,399). Small bowel obstruction is usually caused by adhesions followed by tumors, hernias, foreign bodies, intussusception, gallstones, and inflammatory bowel diseases (400,401). As a rare complication of endometriosis, small bowel obstruction defined as a condition by the presence of endometrial tissue outside of the uterine cavity (402). High fiber diet is appropriate for having the bowels healthy for some people, as fiber, especially insoluble fiber can provide bulk to the stool, which can increase the risk of bowel obstruction. Two doses of modified Dachengqi Decoction herbal formula which was part of traditional Chinese medicine was recommended for patients (403), while Kunitomi *et al.* (404) reported that Daikenchuto did not show significant effectiveness for early bowel obstruction, but it might be suitable for shortening the hospital stay for patients (404). One famous Chinese herbal medicine is Si-Jun-Zi Decoction which composed of four Chinese herbs namely Poria root, licorice root, *Atractylodes macrocephala*, and Ginseng root that can regulate the intestinal immune system, improves the restoration of intestinal function after obstruction by managing intestinal homeostasis which proves its importance for treatment of intestinal obstruction (405).

Anal fistula

The history of anal fistulas goes back to the Ancient Greeks (406,407), which is an epithelized path between anal canal or the rectum and the perianal region (408,409). It is usually challenging to manage the anal fistulas properly (410,411). The anal fistula is the unusual connection between the perianal skin and the anal canal, which can induce to persistent or intermittent purulent drainage (412). It is an internal orifice originating along the anal circumference which is the natural drainage orifice of the infected anal gland (413). Application of a porcine

fistula plug for the treatment of complex anal fistulas is an important and promising technique (414,415). The four types of anal fistula are intersphincteric, transsphincteric, suprasphincteric, and extrasphincter (416-419).

Some of the most important plants for symptomatic treatment of anal fistula are *Zingiber officinale*, *Curcuma longa*, *Commiphora mukul*, *Terminalia chebula*, *Mimosa pudica*, *Embllica officinalis*, *Azadirachta indica*, *Berberis aristata*, *Acacia catechu*, *Cassia tora*, *Leucas cephalotes*, *Rubbia cordifolia*, *Cinnamomum camphora*, *Melia azedarach*, and *Sapindus muskorossi* (420). Santos *et al.* (421) reported that *Baccharis dracunculifolia* extract was able to decrease the lumen of the fistulous tracts and the degree of vascular congestion without totally closing or reducing the local inflammatory process of the fistulous tracts. Comparin *et al.* (422) found that the application of a seton soaked in a glycerin *Aloe barbadensis* Miller extract was important and useful in the treatment of anal fistulas in rats with significantly lower inflammatory process compare to the non-phytotherapeutic seton. The main prescribed herbal components in topical formulation for anal fistula treatment in Persian traditional medicine are gum tragacanth of *Astragalus tragacantha* L., sweat violate oil of *Viola odorata* L., apricot kernel oil of *Prunus armeniaca* L., henna leaf of *Lawsonia alba* Lam., rose oil of *Rosa damascene* Herrm., myrrh (*Commiphora myrrha* (Nees) Engl.), and dry wallflower (*Eryzimum cheiri* (L.) Crantz. (423). Cavasana *et al.* (424) found that the extract of *Aloe vera* was capable to decrease the lumen of the fistulous tract as well as decrease the vascular congestion. Huanglian ointment (HLO) can reduce inflammation and enhance the healing of infected wounds, and its mechanisms are connected to γ -Glutamine and γ -glutamate metabolism, linoleic acid metabolism, tyrosine, phenylalanine, tryptophan biosynthesis, alpha-linolenic acid metabolism, ascorbate, and aldarate metabolism (425). Kanazawa Sutra (KanS) is composed of *Capsicum annuum* tincture, *Ficus carica* latex, and *Achyranthes fauriei* Kshara, and powdered *Curcuma longa* rhizome, can be recommended for treatment of anal fistula (426).

Cyclic vomiting syndrome (CVS)

CVS is clinically known by a pattern of stereotypic, recurrent, and usually incapacitating episodes of intense vomiting and nausea which are identified by periods of time free of signs (427-429). It is known as functional and idiopathic gastrointestinal disorder which related to a disorder of brain-gut interaction appearing in pediatric and adult patients around the world (430-432). It is usually categorized as having four phases: prodromal, acute/vomiting/hyperemesis, recovery, and remission/interepisodic, and the common treatment strategies for it is consisting of trigger avoidance, abortive and prophylactic medication therapies, and supportive care (433). As a disorder of gut-brain interaction (DGBI), it represents 3%-14% of cases of unexplained vomiting and chronic nausea (434,435). It was known as a primarily pediatric illness, which is now identified as the disorder which has a similar prevalence in adults (436,437).

Symptoms and signs of this illness mimic other illnesses such as gastroenteritis, food poisoning, and gastroparesis, which make its recognition difficult or with delayed for some years (438-440). Various types of CVS can be categorized according to the differences in clinical features of episodes, specific triggering contexts, and patterns of associated comorbidities (441,442). It is also reported that CVS possibly has a neurological basis with potential links to neuronal excitability and the mechanism of neural circuits is important for vomiting and nausea (443-445). It has been revealed that CVS are characterized by recurrent attacks of vomiting, nausea, and abdominal pain lasting for minutes in both adolescents and children (446-448). Anxiety is prevalent in pediatric CVS warranting screening, and heart rate variability (HRV) may be used as a biomarker for assessing stress as a trigger for CVS episodes (449,450). Some of the most important medicinal plants and herbs which have been used for many years in South part of Asia for treatment and cure of CVS are *Artemisia maritima* L., *Mentha longifolia* (L.), *Tanacetum faconeri* Hook., *Mentha piperita* L., *Allium cepa* L., *Pimpinella diversifolia* DC., *Hylotelephium telephioides* (Ledeb.)H., *Rheum webbianum* Wall., *Oxyria digyna* (L.), *Viola serpens* Wall., *Persicaria amplexicaulis* (D.Don), *Cichorium intybus* L., *Prunus persica* L., *Ficus carica* L., *Prunus armeniaca* L., *Cucumis sativus* L., *Thymus serpyllum* L., *Persicaria vivipara* (L.), *Saussurea gossypiphora* D., *Rhododendron anthopogon* D., *Aconitum heterophyllum* Wall., *Urtica dioica* L., *Cucurbita maxima*, *Ribes alpestre*, *Rheum austral*, *Morus albla*, *Raphanus sativus* L., *Daucus carota* L., *Oxalis corniculata* L., *Berberis lycium*, *Echinops echinatus* Roxb., *Salvia* sp., *Solanum nigrum* L., *Datura stramonium* L., *Cuscuta reflexa* Roxb., *Swertia petiolate*, *Plantago himalaica*, *Juniperus excelsa*, *Aster vinifera* L., *Rosa indica* L., *Chenopodium album* L., *Morus nigra* L., *Xanthium strumarium* L., *Verbascum Thapsus* L., and *Artemisia annua* L. (451,452). Important treatments for cyclic vomiting syndrome are identify and avoid the triggers, fight signs and symptoms during an episode by using natural relief options for migraines, explore

natural nausea remedies such as chamomile tea, ginger tea, and essential oils such as lemon and peppermint, reduce the fever, minimize diarrhea, and ease dizziness (453-456). Treating cyclic vomiting syndrome can be done through different ways such as medication options such as preventive or prophylactic medications, abortive medications, and lifestyle changes such as avoid cannabis use and address stress (457-462).

Conclusion

Plant-derived components can be applied as synergistic agents and biofilm inhibitors with considering innovative findings using different new technologies for development and effective proof-of-concept research of plant-derived options. Nutraceuticals for gastrointestinal disorders are senna, cascara, *Saccharomyces cerevisiae*, *Andrographis paniculata*, boswellic acids, *Curcuma longa*, purple potato extract, *Aloe vera* gel, alginic acid, anthocyanins, curcumin, quercetin, and resveratrol. The most important medicinal plants for treatment of gallstone are saffron, fenugreek, red radish, cumin, fennel, turmeric, pomegranate, sweet orange and green tea, while the most notable medicinal plants for treatment of gastroesophageal reflux disease are *Curcuma longa*, *Aloe vera*, *Althaea officinalis*, *Morus alba*, *Artemisia asiatica*, and *Fumaria officinalis*. For treatment of peptic ulcer, *Zingiber officinalis*, *Zingiber zerumbet*, *Allium sativum*, *Camellia sinensis*, *Curcuma longa*, and *Artemisia asiatica* are common. Chemical components for treatment of pancreatitis are curcumin, luteolin, ellagic acid, borneol, carvacrol, berberine, emodin, oxymatrine, artesunate, baicalein, baicalin, and calycosin. All in all, in conclusion, this review summarizes the importance of different medicinal plants in treatment of various gastrointestinal diseases. In future, more randomized, large, placebo-controlled researches are needed to evaluate the safety and effectiveness of spices and herbs for treatment of gastrointestinal disorders.

Author Contributions

W.S.: writing-original draft preparations; M.H.S.: writing-original draft preparation, and editing. All authors have read and agreed to the published version of the manuscript.

Funding

Not applicable

Institutional Review Board Statement

Not applicable.

Informed Consent Statement

Not applicable.

Data Availability Statement

Not applicable.

Clinical Trial Number

Not applicable.

Ethics, Consent to Participate, and Consent to Publish Declarations

Not applicable.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Sun W, Shahrajabian MH. Therapeutic potential of phenolic compounds in medicinal plants-natural health products for human health. *Molecules*. 2023;28(1845):1-47. <https://doi.org/10.3390/molecules28041845>
2. Sun W, Shahrajabian MH. The application of arbuscular mycorrhizal fungi as microbial biostimulant, sustainable approaches in modern agriculture. *Plants*. 2023;12(17):3101. <https://doi.org/10.3390/plants12173101>
3. Sun W, Shahrajabian MH, Petropoulos SA, et al. Developing sustainable agriculture systems in medicinal and aromatic plant production by using chitosan and chitin-based biostimulants. *Plants*. 2023;12(13):2469. <https://doi.org/10.3390/plants12132469>
4. Sun W, Shahrajabian MH, Kuang Y, et al. Amino acids biostimulants and protein hydrolysates in agricultural sciences. *Plants*. 2024;13(210):1-22. <https://doi.org/10.3390/plants13020210>
5. Sun W, Shahrajabian MH, Soleymani A. The roles of plant-growth-promoting rhizobacteria (PGPR)-based biostimulants for agricultural production systems. *Plants*. 2024;13(613):1-38. <https://doi.org/10.3390/plants13050613>
6. Shahrajabian MH, Sun W, Soleymani A, et al. Traditional herbal medicines to overcome stress, anxiety and improve mental health in outbreaks of human coronaviruses. *Phytother Res*. 2020;2020(1):1-11. <https://doi.org/10.1002/ptr.6888>
7. Shahrajabian MH, Sun W, Cheng Q. Traditional herbal medicine for the prevention and treatment of cold and flu in the autumn of 2020, overlapped with Covid-19. *Nat Prod Commun*. 2020;15(8):1-10. <https://doi.org/10.1177/1934578X20951431>
8. Shahrajabian MH, Sun W, Shen H, et al. Chinese herbal medicine for SARS and SARS-CoV-2 treatment and prevention, encouraging using herbal medicine for Covid-19 outbreak. *Acta Agric Scand Sect B Soil Plant Sci*. 2020;70(5):437-443. <https://doi.org/10.1080/09064710.2020.1763448>
9. Shahrajabian MH. Medicinal herbs with anti-inflammatory activities for natural and organic healing. *Curr Org Chem*. 2021;25(23):1-17. <https://doi.org/10.2174/138527282566621110115656>
10. Shahrajabian MH, Sun W, Cheng Q. Different methods for molecular and rapid detection of human novel coronavirus. *Curr Pharm Des*. 2021;27:1-10. <https://doi.org/10.2174/1381612827666210604114411>
11. Machate DJ, Figueiredo PS, Marcelino G, et al. Fatty acid diets: regulation of gut microbiota composition and obesity and its related metabolic dysbiosis. *Int J Mol Sci*. 2020;21(11):4093. <https://doi.org/10.3390/ijms21114093>
12. Filippis AD, Ullah H, Baldi A, et al. Gastrointestinal disorders and metabolic syndrome: Dysbiosis as a key link and common bioactive dietary components useful for their treatment. *Int J Mol Sci*. 2020;21(14):4929. <https://doi.org/10.3390/ijms21144929>
13. Shahrajabian MH, Chaski C, Polyzos N, et al. Biostimulants application: A low input cropping management tool for sustainable farming of vegetables. *Biomolecules*. 2021;11(5):698. <https://doi.org/10.3390/biom11050698>
14. Shahrajabian MH, Sun W. Using sumac (*Rhus coriaria* L.), as a miraculous spice with outstanding pharmacological activities. *Not Sci Biol*. 2022;14(1):1-14. <https://doi.org/10.15835/nsb14111118>
15. Parisio C, Lucarini E, Micheli L, et al. Pomegranate mesocarp against colitis-induced visceral pain in rats: Effects of a decoction and its fractions. *Int J Mol Sci*. 2020;21(12):4304. <https://doi.org/10.3390/ijms21124304>
16. Gao X, Liu J, Li L, et al. A brief review of nutraceutical ingredients in gastrointestinal disorders: Evidence and suggestions. *Int J Mol Sci*. 2020;21(5):1822. <https://doi.org/10.3390/ijms21051822>
17. Martinez V, De-Hond AI, Borrelli F, et al. Cannabidiol and other non-psychoactive cannabinoids for prevention and treatment of gastrointestinal disorders: useful nutraceuticals? *Int J Mol Sci*. 2020;21(9):3067. <https://doi.org/10.3390/ijms21093067>
18. Embeya VO, Simbi J-BL, Stevigny C, et al. Traditional plant-based remedies to control gastrointestinal disorders in livestock in the regions of Kamina and Kaniama (Katanga province, Democratic Republic of Congo). *J Ethnopharmacol*. 2014;153(3):686-693. <https://doi.org/10.1016/j.jep.2014.03.027>
19. Colosimo R, Warren FJ, Edwards CH, et al. Comparison of the behavior of fungal and plant cell wall during gastrointestinal digestion and resulting health effects: A review. *Trends Food Sci Technol*. 2021;110:132-141. <https://doi.org/10.1016/j.tifs.2021.02.001>
20. Rokaya MB, Uprety Y, Poudel RC, et al. Traditional uses of medicinal plants in gastrointestinal disorders in Nepal. *J Ethnopharmacol*. 2014;158:221-229. <https://doi.org/10.1016/j.jep.2014.10.014>
21. Thumann TA, Pferschy-Wenzig E-M, Moissl-Eichinger C, et al. The role of gut microbiota for the activity of medicinal plants traditionally used in the European Union for gastrointestinal disorders. *J Ethnopharmacol*. 2019;245:112153. <https://doi.org/10.1016/j.jep.2019.112153>

22. Guo D, Zhang W, Yang H, et al. Celastrol induces necroptosis and ameliorates inflammation via targeting biglycan in human gastric carcinoma. *Int J Mol Sci.* 2019;20(22):5716. <https://doi.org/10.3390/ijms20225716>
23. Luz BBD, Maria-Ferreira D, Dallazen JL, et al. Effectiveness of the polyphenols-rich *Sedum dendroideum* infusion on gastric ulcer healing in rats: Roles of protective endogenous factors and antioxidant and anti-inflammatory mechanisms. *J Ethnopharmacol.* 2021;278:114260. <https://doi.org/10.1016/j.jep.2021.114260>
24. Ahmad S, Ullah F, Ayaz M, et al. Antioxidant and anticholinesterase investigations of *Rumex hastatus* D. Don: potential effectiveness in oxidative stress and neurological disorders. *Biol Res.* 2015;48(1):1-8. <https://doi.org/10.1184/s40659-015-0010-2>
25. Benvenuti DF, Monache FD, Filho VC, et al. Phytochemical analysis and gastroprotective activity of the root bark from *Maytenus robusta*. *Nat Prod Commun.* 2016;11(5):597-599.
26. Gupta M, Kapoor B, Gupta R, et al. Plants and phytochemicals for treatment of peptic ulcer: An overview. *South Afr J Bot.* 2021;138:105-114. <https://doi.org/10.1016/j.sajb.2020.11.030>
27. Kumar MR, Niyas M, Mani TT, et al. A review on medicinal plants for peptic ulcer. *Der Pharm Lett.* 2011;3(2):180-186.
28. Mitra PK. Comparative evaluation of anti-gastric ulcer activity of root, stem, and leaves of *Amaranthus spinosus* Linn. in rats. *Int J Herb Med.* 2013;1(2):22-29.
29. Prayoga DK, Aulifa DL, Budiman A, et al. Plants with anti-ulcer activity and mechanism: A review of preclinical and clinical studies. *Drug Des Develop Ther.* 2024;2024:193-213. <https://doi.org/10.2147/DDDT.S446949>
30. Shatri AMN, Mumbengegwi DR. Ethnomedicinal use and phytochemical analysis of medicinal plants used to treat gastrointestinal conditions by Awambo people in Iikokola village, Namibia *Sci Afr.* 2022;18:e01428. <https://doi.org/10.1016/j.sciaf.2022.e01428>
31. Sulaiman AN, Arzai AH, Taura DW. Ethnobotanical survey: A comprehensive review of medicinal plants used in treatment of gastro intestinal diseases in Kano state, Nigeria. *Phytomed Plus.* 2022;2(1):100180. <https://doi.org/10.1016/j.phyplu.2021.100180>
32. Shane AL, Mody RK, Crump JA, et al. Infectious diseases society of America clinical practice guidelines for the diagnosis and management of infectious diarrhea. *Clin Infect Dis.* 2017;65:e45-e80.
33. Shahrajabian MH, Sun W. Mechanism of action of collagen and epidermal growth factor: A review on theory and research methods. *Mini Rev. Med Chem.* 2024;24(4):453-477. <https://doi.org/10.2174/1389557523666230816090054>
34. Shahrajabian MH, Sun W. The significance and importance of dPCR, qPCR, and SYBR green PCR kit in the detection of numerous diseases. *Curr Pharm Des.* 2024;30(3):1-11. <https://doi.org/10.2174/0113816128276560231218090436>
35. Shahrajabian MH, Sun W. Multidimensional uses of bitter melon (*Momordica charantia* L.) considering the important functions of its chemical components. *Curr Org Synth.* 2024. <https://doi.org/10.2174/0115701794285586240523101245>
36. Ranfaing J, Dunyach-Remy C, Lavigne J-P, et al. Propolis potentiates the effect of cranberry (*Vaccinium macrocarpon*) in reducing the motility and the biofilm formation of uropathogenic *Escherichia coli*. *Plos One.* 2018;13:e0202609.
37. Chen JC, Huang LJ, Wu SL, et al. Ginger and its bioactive component inhibit enterotoxigenic *Escherichia coli* heat-labile enterotoxin-induced diarrhea in mice. *J Agric Food Chem.* 2007;55:8390-8397.
38. Sugita-Konishi Y, Hara-Kudo Y, Amano F, et al. Epigallocatechin gallate and gallic acid gallate in green tea catechins inhibit extracellular release of Vero toxin from enterohemorrhagic *Escherichia coli* O157:H7. *Biochim Biophys Acta.* 1999;1472:42-50.
39. Dadi PK, Ahmad M, Ahmad Z. Inhibition of ATPase activity of *Escherichia coli* ATP synthase by polyphenols. *Int J Biol Macromol.* 2009;45:72-79.
40. Garg SS, Gupta J, Sharma S, et al. An insight into the therapeutic applications of coumarin compounds and their mechanisms of action. *Eur J Pharm. Sci.* 2020;152:105424.
41. Sack RB, Froehlich JL. Berberine inhibits intestinal secretory response of *Vibrio cholerae* and *Escherichia coli* enterotoxins. *Infect Immun.* 1982;35:471-475.
42. Amin AH, Subbaiah TV, Abbasi KM. Berberine sulfate: Antimicrobial activity, bioassay, and mode of action. *Can J Microbiol.* 1969;15:1067-1076.
43. Baskaran SA, Kollanoor-Johny A, Nair MS, et al. Efficacy of plant-derived antimicrobials in controlling enterohemorrhagic *Escherichia coli* virulence in vitro. *J Food Prot.* 2016;79:1965-1970.

44. Mooyottu S, Kollanoor-Johny A, Flock G, et al. Carvacol and trans-cinnamaldehyde reduce *Clostridium difficile* toxin production and cytotoxicity in vitro. *Int J Mol Sci.* 2014;15:4415-4430.
45. Thapa D, Louis P, Losa R, et al. Essential oils have different effects on human pathogenic and commensal bacteria in mixed faecal fermentations compared with pure cultures. *Microbiology.* 2015;161:441-449.
46. Zhang Y, Qi Z, Liu Y, et al. Baicalin protects mice from lethal infection by Enterohemorrhagic *Escherichia coli*. *Front Microbiol.* 2017;8:395.
47. Kemal ME, Bakchiche B, Kemal M, et al. Six Algerian plants: Phenolic profile, antioxidant, antimicrobial activities associated with different simulated gastrointestinal digestion phases and antiproliferative properties. *J Herb Med.* 2023;38:100636. <https://doi.org/10.1016/j.hermed.2023.100636>
48. Lauche R, Kumar S, Hallmann J, et al. Efficacy and safety of Ayurvedic herbs in diarrhoea-predominant irritable bowel syndrome: A randomized controlled crossover trial. *Complement Ther Med.* 2016;26:171-177.
49. Tal-Dia A, Toure K, Sarr O, et al. A baobab solution for the prevention and treatment of acute dehydration in infantile diarrhea. *Dakar Med.* 1997;42:68-73.
50. Chauhan RK, Jain AM, Dube MK, et al. A combination of sulfadimidine, neomycin, and berberine in the treatment of infections diarrhea. *Indian J Pediatr.* 1969;36:242-244.
51. Zhou FF, Wu S, Klena JD, et al. Clinical characteristics of *Clostridium difficile* infection in hospitalized patients with antibiotic-associated diarrhea in a university hospital in China. *Eur J Clin Microbiol Infect Dis.* 2014;33:1773-1779.
52. Khin Maung U, Myo K, et al. Clinical trial of berberine in acute watery diarrhoea. *Br Med J Clin Res Ed.* 1985;291:1601-1605.
53. DuPont HL, Flores Sanchez J, Ericsson CD, et al. Comparative efficacy of loperamide hydrochloride and bismuth subsalicylate in the management of acute diarrhea. *Am J Med.* 1990;88:15s-19s.
54. Khan N, Ahmed M, Ahmed A, et al. Important medicinal plants of chitral gol National park (cgnp) Pakistan. *Pakistan J Bot.* 2011;43(2):797-809.
55. Dib I, Angenot L, Mihamou A, et al. *Artemisia campestris* L.: Ethnomedicinal, phytochemical and pharmacological review. *J Herbal Med.* 2017;7:1-10.
56. Khan SW, Abbas Q, Hassan SN, et al. Medicinal plants of Turmic valley (Central Karakoram National Park), Gilgit-Baltistan, Pakistan. *J Bioresour Manage.* 2015;2(2):11.
57. Rahimi Y, Taleei A, Ranjbar M. Long-term water deficit modulates antioxidant capacity of peppermint (*Mentha piperita* L.). *Sci Hortic.* 2018;237:36-43.
58. Mahendra G, Rahman LU. Ethnomedicinal, phytochemical and pharmacological updates on peppermint (*Mentha x piperita* L.)- A review. *Phytother Res.* 2020;34(9):2088-2139. <https://doi.org/10.1002/ptr.6664>
59. Karunakaran G, Jagathambal M, Kumar GS, et al. *Hylotelephium telephium* flower extract-mediated biosynthesis of CuO and ZnO nanoparticles with promising antioxidant and antibacterial properties for healthcare applications. *JOM.* 2020;72(3):1264-1272.
60. Kumar A, Chauhan P, Bhardwaj V, et al. In vitro antioxidant and phytochemical investigations of ethanolic extracts of *Viola serpens* and *Morus nigra*. *J Chem Pharm Res.* 2011;3(4):166-171.
61. Kumar P, Digvijay S. Assessment of genetic diversity of *Viola serpens* Wall. in Himachal Pradesh using molecular markers. *World J Pharm. Res.* 2014;3:2716-2726.
62. Batool S, Ahmed MS. *Bistorta amplexicaulis*: A brief insight to its ethnobotany. *J Bioresour Manag.* 2016;3(3):2.
63. Mishra AP, Saklani S, Parcha V, et al. Antibacterial activity and phytochemical characterization of *Saussurea gossypiphora* D. Don. *Arch Microbiol.* 2021; 2021:1-11. <https://doi.org/10.1007/s00203-021-02494-1>
64. Paramanick D, Panday R, Shukla SS, et al. Primary pharmacological and other important findings on the medicinal plant *Aconitum heterophyllum* (aruna). *J Pharmacopunct.* 2017;20(2):89. <https://doi.org/10.3831/KPL.2017.20.011>
65. Maurya SK, Kushwaha AK, Seth A. Ethnomedicinal review of Usnakantaka (*Echinops echinatus* Roxb.). *Pharmacogn Rev.* 2015;9(18):149. <https://doi.org/10.4103/0973-7847.162138>
66. Ahmad AA, Kasim KF, MaRadzi AH, et al. Peptic ulcer: Current prospects of diagnostic and nanobiotechnological trends on pathogenicity. *Process Biochem.* 2019;85:51-59. <https://doi.org/10.1016/j.procbio.2019.06.024>
67. Singh G, Rawat G. Ethnomedicinal survey of Kedarnath wildlife sanctuary in Western Himalaya, India. *Indian J Fundament Appl Life Sci.* 2011;1(1):35-46.
68. Sargin SA, Selvi S, Lopez V. Ethnomedicinal plants of Sarigol district (Manisa), Turkey. *J Ethnopharmacol.* 2015;171:64-84. <https://doi.org/10.1016/j.jep.2015.05.031>
69. Dawa S, Gurmet P, Stobgais T, et al. Survey and ethno-botanical study of medicinal plants of some selected villages of Singay-Lalok region of Leh (UT Ladakh Region). *Asian J Res Bot.* 2021; 2021:60-75.

70. Rahnavard R, Razavi N. A review on the medical effects of *Capparis spinosa* L. *Adv Herbal Med.* 2017;3(1):44-53.
71. Zhang S, Lin L, Liu W, et al. Shen-Ling-Bai-Zhu-San alleviates functional dyspepsia in rats and modulates the composition of the gut microbiota. *Nutr Res.* 2019;71:89-99. <https://doi.org/10.1016/j.nutres.2019.10.001>
72. Sadiq A, Hayat MQ, Ashraf M. Ethnopharmacology of *Artemisia annua* L.: A review. *Artemisia annua. Pharmacol Biotechnol.* 2017; 2017:9-25.
73. Hadi A, Singh S. Ethno-botanical studies of some threatened medicinal plants and local perception of its population decline in Kargil, Ladakh UT. *J Med Plants.* 2021;9(4):44-50.
74. Jishtu V, Bhondge SW, Bhushan B, et al. Threatened ethnomedicinal plants of Dodra-Kwar region of Himachal Pradesh, NW Himalaya. *J Med Plants.* 2021;9(2):151-159.
75. Phromnoi K, Sinchaiyakij P, Khanaree C, et al. Anti-inflammatory and antioxidant activities of medicinal plants used by traditional healers for antiulcer treatment. *Sci Pharm.* 2019;87:1-22. <https://doi.org/10.3390/scipharm87030022>
76. Direito R, Barbalho SM, Figueira ME, et al. Medicinal plants, phytochemicals and regulation of the NLRP3 inflammasome in inflammatory bowel diseases: A comprehensive review. *Metabolites.* 2023;13(6):728. <https://doi.org/10.3390/metabo13060728>
77. Sakunpak A, Panichayupakaranant P. Antibacterial activity of Thai edible plants against gastrointestinal pathogenic bacteria and isolation of a new broad spectrum antibacterial polyisoprenylated benzophenone, chamuangone. *Food Chem.* 2012;130(4):826-831. <https://doi.org/10.1016/j.foodchem.2011.07.088>
78. Darvishpor S, Hosseini A, Davoodi A, et al. A review on medicinal plants used for nausea and vomiting in Persian medicine. *Global J Med Res Pharma Drug Discov Toxicol Med.* 2018;18(1):1-18.
79. Shahrajabian MH, Sun W. Medicinal plants, economical and natural agents with antioxidant activity. *Curr Nutr Food Sci.* 2022;19(8):763-784. <https://doi.org/10.2174/1573401318666221003110058>
80. Shahrajabian MH, Sun W. Sustainable approaches to boost yield and chemical constituents of aromatic and medicinal plants by application of biostimulants. *Recent Pat Food Nutr Agric.* 2022;13(2):72-92. <https://doi.org/10.2174/2772574X13666221004151822>
81. Shahrajabian MH, Sun W. Asparagus (*Asparagus officinalis* L.) and pennyroyal (*Mentha pulegium* L.), impressive advantages with wondrous health-beneficial phytochemicals. *Not Sci Biol.* 2022;14(2):11212. <https://doi.org/10.55779/nsb14211212>
82. Shahrajabian MH, Sun W, Cheng Q. The importance of flavonoids and phytochemicals of medicinal plants with antiviral activities. *Mini Rev Org Chem.* 2022;19(3):293-318. <https://doi.org/10.2174/1570178666210707161025>
83. Shahrajabian MH, Cheng Q, Sun W. The organic life according to traditional Chinese medicine with anti-cancer approaches. *Curr Nutr Food Sci.* 2022;18(8):1-6. <https://doi.org/10.2174/1871520622666220425093907>
84. Elmaghraby DA, Alsalman GA, Alawadh LH, et al. Integrated traditional herbal medicine in the treatment of gastrointestinal disorder: the pattern of use and the knowledge of safety among the Eastern region Saudi population. *BMC Complement Med Ther.* 2023;23:373. <https://doi.org/10.1186/s12906-023-04197-0>
85. Laurindo LF, Santos ARDOD, Carvalho ACAD, et al. Phytochemicals and regulation of NF- κ B in inflammatory Bowel diseases: An overview of in vitro and in vivo effects. *Metabolites.* 2023;13(1):96. <https://doi.org/10.3390/metabo13010096>
86. Alkhatib M, Fayad C, Badran A, et al. Preventive therapeutics effects of *Punica granatum* (Pomegranate) in respiratory and digestive diseases: A review. *Appl Sci.* 2022;12(23):12326. <https://doi.org/10.3390/app122312326>
87. Qazi NG, Khan A-U, Abbasi SW, et al. Inhibition and PDR mediated signaling: Toxicology evaluation on vital organs. *Molecules.* 2022;27(18):5919. <https://doi.org/10.3390/molecules27185919>
88. Li L, Liu C, Xia T, et al. Association between plant-based dietary index gallstone disease: A cross sectional study from NHANES. *Plos One.* 2024;19(6):e0305822. <https://doi.org/10.1371/journal.pone.0305822>
89. Jiang Z-Y, Jiang C-Y, Wang L, et al. Increase NPC1L1 and ACAT2 expression in the jejunal mucosa from Chinese gallstone patients. *Biochem Biophys Res Commun.* 2009;379(1):49-54. <https://doi.org/10.1016/j.bbrc.2008.11.131>
90. Shahrajabian MH, Sun W, Cheng Q. Using bacteria and fungi as plant biostimulants for sustainable agricultural production systems. *Recent Pat Biotechnol.* 2022;17(3):206-244. <https://doi.org/10.2174/1872208316666220513093021>
91. Shahrajabian MH, Sun W. The importance of traditional Chinese medicine in the intervention and treatment of HIV while considering its safety and efficacy. *Curr HIV Res.* 2023;21(6):1-16. <https://doi.org/10.2174/011570162X271199231128092621>

92. Shahrajabian MH, Sun W. Iranian traditional medicine (ITM) and natural remedies for treatment of the common cold and flu. *Rev Recent Clin Trials*. 2023;19(2):91-100. <https://doi.org/10.2174/0115748871275500231127065053>
93. Shahrajabian MH, Sun W. Survey on multi-omics, and multi-omics data analysis, integration and application. *Curr Pharm Anal*. 2023;19(4):267-281. <https://doi.org/10.2174/1573412919666230406100948>
94. Nissinen MJ, Pitkanen N, Simonen P, et al. Genetic polymorphism of sterol transporters in children with future gallstones. *Digest Liver Dis*. 2018;50(9):954-960. <https://doi.org/10.1016/j.dld.2018.04.007>
95. Ding L, Wang S, Jiang W, et al. Identification of intestinal microbial community in gallstone patients with metagenomic next-generation sequencing. *Diagnostics*. 2023;13(16):2712. <https://doi.org/10.3390/diagnostics13162712>
96. Shabanzade DM. The symptomatic outcomes of cholecystectomy for gallstones. *J Clin Med*. 2023;12(5):1897. <https://doi.org/10.3390/jcm12051897>
97. Zhang D, Wu C, Liu Y, et al. Microplastics are detected in human gallstones and have the ability to form large cholesterol-microplastic heteroaggregates. *J Hazard Mater*. 2024;467:133631. <https://doi.org/10.1016/j.hazmat.2024.133631>
98. Mendez-Sanchez N, Zamora-Valdes D, Chavez-Tapia NC, et al. Role of diet in cholesterol gallstone formation. *Clin Chim Acta*. 2007;376(1-2):1-8. <https://doi.org/10.1016/j.cca.2006.08.036>
99. Wirth J, Joshi AD, Song M, et al. A healthy lifestyle pattern and the risk of symptomatic gallstone disease: results from 2 prospective cohort studies. *Am J Clin Nutr*. 2020;112(3):586-594. <https://doi.org/10.1093/ajcn.nqaa154>
100. Xia S, Wang J, Kalionis B, et al. Genistein protects against acute pancreatitis via activation of an apoptotic pathway mediated through endoplasmic reticulum stress in rats. *Biochem Biophys Res Commun*. 2019;509(2):421-428. <https://doi.org/10.1016/j.bbrc.2018.12.108>
101. Arrout A, Ghallab YE, Elmakssoudi A, et al. Black cumin and clove: Litholytic volatile compounds and inhibitors of inflammation-induced gallstone. *Sci Afr*. 2024;23:e02110. <https://doi.org/10.1016/j.sciaf.2024.e02110>
102. Arrout A, Ghallab YE, Yafout M, et al. Medicinal plants for gallstones: A cross-sectional survey of Moroccan patients. *Phytomed Plus*. 2024;4(1):100524. <https://doi.org/10.1016/j.phyplu.2024.100524>
103. Chang, C.-M.; Chiu THT, Chang C-C, et al. Plant-based diet, cholesterol, and risk of gallstone disease: A prospective study. *Nutrients*. 2019;11:335. <https://doi.org/10.3390/nu11020335>
104. Cheng J, Ouwehand AC. Gastroesophageal reflux disease and probiotics: A systematic review. *Nutrient*. 2020;12(1):132. <https://doi.org/10.3390/nu12010132>
105. Hosseini M, Salari M, Salari R. Psyllium seed may be effective in the treatment of gastroesophageal reflux disease (GERD) in patients with functional constipation. *J Med Hypth Ideas*. 2015;9(2):S4-S7. <https://doi.org/10.1016/j.jmhi.2015.11.002>
106. Pados BF, Davitt ES. Pathophysiology of gastroesophageal reflux disease in infants and nonpharmacologic strategies for symptom management. *Nurs Women Health*. 2020;24(2):101-114. <https://doi.org/10.1016/j.nwh.2020.01.005>
107. Ferguson DD, DeVault KR. Medical management of gastroesophageal reflux disease. *Expert Opin Pharmacother*. 2007;8(1):39-47. <https://doi.org/10.1517/14656566.8.1.39>
108. Chapelle N, Ghezala IB, Barkun A, et al. The pharmacotherapeutic management of gastroesophageal reflux disease (GERD). *Expert Opin Pharmacother*. 2021;22(2):219-227. <https://doi.org/10.1080/14656566.2020.1817385>
109. Shahrajabian MH, Sun W. Various techniques for molecular and rapid detection of infectious and epidemic diseases. *Lett Org Chem*. 2023;20(9):779-801. <https://doi.org/10.2174/1570178620666230331095720>
110. Shahrajabian MH, Sun W. Study of different types of fermentation in wine-making process and considering aromatic substances and organic acid. *Curr Org Synth*. 2023;20. <https://doi.org/10.2174/1570179420666230803102253>
111. Shahrajabian MH, Petropoulos SA, Sun W. Survey of the influences of microbial biostimulants on horticultural crops: Case studies and successful paradigms. *Horticulturae*. 2023;9(193):1-24. <https://doi.org/10.3390/horticulturae9020193>
112. Shahrajabian MH, Sun W. Assessment of wine quality, traceability and detection of grapes wine, detection of harmful substances in alcohol and liquor composition analysis. *Lett Drug Des Discov*. 2024;21(8):1377-1399. <https://doi.org/10.2174/1570180820666230228115450>

113. Shahrajabian MH, Sun W. The golden spice for life: Turmeric with the pharmacological benefits of curcuminoids components, including curcumin, bisdemethoxycurcumin, and demethoxycurcumins. *Curr Org Synth.* 2024;21(5):665-683. <https://doi.org/10.2174/1570179420666230607124949>
114. Shahrajabian MH, Sun W. The power of the underutilized and neglected medicinal plants and herbs of the Middle East. *Rev Recent Clin Trials.* 2024. <https://doi.org/10.2174/011574881276544240212105612>
115. Shahrajabian MH, Sun W. Importance of thymoquinone, sulforaphane, phloretin, and epigallocatechin and their health benefits. *Lett Drug Des Discov.* 2024;21(2):209-225. <https://doi.org/10.2174/1570180819666220902115521>
116. Bhardwaj K, Kishore L. Natural remedies: For gastroesophageal reflux disease. *J Med Plants Stud.* 2021;9(4):114-118.
117. Rizzo G, Baroni L, Bonetto C, et al. The tole of a plant-only (vegan) diet in gastroesophageal reflux disease: Online survey of the Italian general population. *Nutrients.* 2023;15(22):4725. <https://doi.org/10.3390/nu15224725>
118. Herdiana Y. Functional food in relation to gastroesophageal reflux disease (GERD). *Nutrients.* 2023;15:3583. <https://doi.org/10.3390/nu15163583>
119. Shih Y-S, Tsai C-H, Li T-C, et al. Effect of wu chu yu tang on gastroesophageal reflux disease: Randomized, double-blind, placebo controlled trial. *Phytomedicine.* 2019;56:118-125. <https://doi.org/10.1016/j.phymed.2018.09.185>
120. Mehraban MSA, Shirzad M, Ahmadian-Attari M, et al. Effect of rose oil on gastroesophageal reflux disease in comparison with omeprazole: A double-blind controlled trial. *Complement. Ther. Clin. Pract.* 2021;43:101361. <https://doi.org/10.1016/j.ctcp.2021.101361>
121. Zohalinezhad ME, Imanieh MH, Samani SM, et al. Effect of quince syrup on clinical symptoms of children with symptomatic gastroesophageal reflux disease: A double-blind randomized controlled clinical trial. *Complement Ther Clin Pract.* 2015;21(4):268-276. <https://doi.org/10.1016/j.ctcp.2015.09.005>
122. Amodio E, Grazia SD, Genovese D, et al. Clinical and epidemiologic features of viral gastroenteritis in hospitalized children: An 11-year surveillance in Palermo (Sicily). *Viruses* 2023;15(1):41. <https://doi.org/10.3390/v15010041>
123. Chong KC, Chan EYY, Lee TC, et al. A 21-year retrospective analysis of environmental impacts on paediatric acute gastroenteritis in an affluent setting. *Sci Total Environ.* 2021;764:142845. <https://doi.org/10.1016/j.scitotenv.2020.142845>
124. Adedire O, Love NK, Hughes HE, et al. Early detection and monitoring of gastrointestinal infections using syndromic surveillance: A systematic review. *Int. J. Environ. Res. Public Health.* 2024;21(4):489. <https://doi.org/10.3390/ijerph21040489>
125. Hasan H, Nasirudeen NA, Ruzlan MAF, et al. Acute infectious gastroenteritis: The causative agents, omics-based detection of antigens and novel biomarkers. *Children.* 2021;8(12):1112. <https://doi.org/10.3390/children8121112>
126. Mitra AK, Asala AF, Malone S, et al. Effects of probiotics in adults with gastroenteritis: A systematic review and meta-analysis of clinical trials. *Disease.* 2023;11(4):138. <https://doi.org/10.3390/diseases11040138>
127. Duan X, Li H, Tan X, et al. Polygonum cillinerve polysaccharide inhibits transmissible gastroenteritis virus by regulating microRNA-181. *The Vet J.* 2024;304:106083. <https://doi.org/10.1016/j.tvjl.2024.106083>
128. Tan X, Cui J, Liu N, et al. Study on the immune-enhancing and inhabiting transmissible gastroenteritis virus effects of polysaccharides from Cimicifuga rhizoma. *Microb. Pathogen.* 2024;192:106719. <https://doi.org/10.1016/j.micpath.2024.106719>
129. Shetty YC, Koli PG, Lahoti M, et al. A prospective, single centre, open label, single arm pilot study to evaluate the efficacy and safety of Amlapitta Mishran Suspension in participants with endoscopic gastritis. *J Ayurveda Integr Med.* 2022;13(4):100664. <https://doi.org/10.1016/j.jaim.2022.100664>
130. Wang R, Zhao Y, Zhou L, et al. Costunolide ameliorates MNNG-induced chronic atrophic gastritis through inhibiting oxidative stress and DNA damage via activation of Nrf2. *Phytomedicine.* 2024;130:155581. <https://doi.org/10.1016/j.phymed.2024.155581>
131. Sangiovanni E, Piazza S, Vrhovsek U, et al. A bio-guided approach for the development of a chestnut-based proanthocyanidin-enriched nutraceutical with potential anti-gastritis properties. *Pharmacol Res.* 2018;134:145-155. <https://doi.org/10.1016/j.phrs.2018.06.016>
132. Young E, Edwards L, Singh R. The role of artificial intelligence in colorectal cancer screening: Lesion detection and lesion characterization. *Cancers.* 2023;15(21):5126. <https://doi.org/10.3390/cancers15215126>

133. Luo H, Liu L, Zou J, et al. Network pharmacology study of *Scleromitrion diffusa* (Willd). *R. J. Wang and Scutellaria barbata* D. Don in the treatment of chronic atrophic gastritis. *Intell Pharm.* 2024;2(1):45-50. <https://doi.org/10.1016/j.ipha.2023.10.003>
134. Lee J-H, Lee D-U, Jeong C-S. *Gardenia jasminoides* Ellis ethanol extract and its constituents reduce the risks of gastritis and reverse gastric lesions in rats. *Food Chem Toxicol.* 2009;47(6):1127-1131. <https://doi.org/10.1016/j.fct.2009.01.037>
135. Wafaey AA, El-Hawary SS, Abdelhameed MF, et al. Green synthesis of zinc oxide nanoparticles using ethanolic extract of *Gliricidia sepium* (Jacq.) Kunth. Ex. Walk., stem: Characterization and their gastroprotective effect on ethanol-induced gastritis in rats. *Bioorg Chem.* 2024;145:107225. <https://doi.org/10.1016/j.bioorg.2024.107225>
136. Hu C, Cao F, Jiang Y, et al. Molecular insights into chronic atrophic gastritis treatment: *Coptis chinensis* Franch studied via network pharmacology, molecular dynamics simulation and experimental analysis. *Comput Biol Med.* 2024;178:108804. <https://doi.org/10.1016/j.compbimed.2024.108804>
137. Song J, Xu X, He S, et al. Identification of the therapeutic effect and molecular mechanism of *Coptis chinensis* Franch. and *Magnolia officinalis* var. *biloba* on chronic gastritis. *J Ethnopharmacol.* 2023;317:116864. <https://doi.org/10.1016/j.jep.2023.116864>
138. Lou N, Zhai M, Su Z, et al. Pharmacodynamics and pharmacological mechanism of Moluodan concentrated pill in the treatment of atrophic gastritis: A network pharmacological study and in vivo experiments. *J Ethnopharmacol.* 2024;318:116937. <https://doi.org/10.1016/j.jep.2023.116937>
139. Liu Z, Shang M, Wang Z, et al. Studies on the active constituents and the related mechanisms of Jinhong Tablet in the treatment of chronic gastritis based on network pharmacology and in vivo experimental validation. *Pharmacol. Res. Modern Chin Med.* 2023;7:100247. <https://doi.org/10.1016/j.prmcm.2023.100247>
140. Yang X, Wang J, Guo J, et al. Therapeutic effect of ginger on gastritis: Regulation of STAT3/MAPK signaling pathway and gastrointestinal hormone balance. *J Funct Foods.* 2024;119:106279. <https://doi.org/10.1016/j.jff.2024.106279>
141. Nam HH, Choo BK. *Geranium koreanum*, a medicinal plant *Geranii Herba*, ameliorate the gastric mucosal injury in gastritis-induced mice. *J Ethnopharmacol.* 2021;265:113041. <https://doi.org/10.1016/j.jep.2020.113041>
142. Zhang R, Qin X, Liu, Y. Exploration of the intestinal flora to reveal the important contribution of *Radix Astragali* to *Huangqi Jianzhong Tang* in treating chronic atrophic gastritis rats. *J Pharm Biomed Anal.* 2024;242:116067. <https://doi.org/10.1016/j.jpba.2024.116067>
143. Boeing T, Silva LMD, Somensi LB, et al. Antiulcer mechanisms of *Vernonia condensata* Baker: A medicinal plant used in the treatment of gastritis and gastric ulcer. *J Ethnopharmacol.* 2016;184:196-207. <https://doi.org/10.1016/j.jep.2016.02.049>
144. Dhondrup R, Tidwell T, Zhang X, et al. Tibetan medicine Liuwei Muxiang pills (LWMX pills) effectively protects mice from chronic non-atrophic gastritis. *Phytomedicine.* 2023;115:154826. <https://doi.org/10.1016/j.phymed.2023.154826>
145. Liu Y, Yuang Y-Y, Geng Z-P, Li J-L, et al. Elucidation of the potential molecular mechanism of the active compounds of *Bryophyllum pinnatum* (L. F.) Oken against gastritis based on network pharmacology. *Chin J Anal Chem.* 2023;51(1):100193. <https://doi.org/10.1016/j.cjac.2022.100193>
146. Chen M, Li Y, Li L, et al. Qi-Zhi-Wei-Tong granules alleviates chronic non-atrophic gastritis in mice by altering the gut microbiota and bile acid metabolism. *J. Ethnopharmacol.* 2024, 319, 117304. <https://doi.org/10.1016/j.jep.2023.117304>
147. Chen H-F, Gong Y, Huang Z, et al. Efficacy and safety of Chinese herbal medicine Qirui Weishu capsule in treating chronic non-atrophic gastritis: A multicentre, double-blind, randomized controlled clinical trial. *J Ethnopharmacol.* 2022;294:115341. <https://doi.org/10.1016/j.jep.2022.115341>
148. Han L, Li T, Wang Y, et al. Weierning, a Chinese patent medicine, improves chronic atrophic gastritis with intestinal metaplasia. *J Ethnopharmacol.* 2023;309:116345. <https://doi.org/10.1016/j.jep.2023.116345>
149. Tong Y, Wang R, Liu X, et al. Zuojin Pill ameliorates chronic atrophic gastritis induced by MNNG through TGF-B1/PI3L/Akt axis. *J Ethnopharmacol.* 2021;271:113893. <https://doi.org/10.1016/j.jep.2021.113893>
150. Qin F, Liu J-Y, Yuan J-H. Chaihu-Shugan-San, an oriental herbal preparation, for the treatment of chronic gastritis: A meta-analysis of randomized controlled trials. *J Ethnopharmacol.* 2013;146(2):433-439. <https://doi.org/10.1016/j.jep.2013.01.029>

151. Kim HG, Kim M-Y, Cho JY. *Alisma canaliculatum* ethanol extract suppresses inflammatory responses in LPS-stimulated macrophages, HC/EtOH-induced gastritis, and DSS-triggered colitis by targeting Src/Syk and TAK1 activities. *J Ethnopharmacol.* 2018;219:202-212. <https://doi.org/10.1016/j.jep.2018.03.022>
152. Shin JH, Lee CW, Oh SJ, et al. Protective effect of silymarin against ethanol-induced gastritis in rats: Role of sulfhydryls, nitric oxide and gastric sensory afferents. *Food Chem Toxicol.* 2013;55:353-357. <https://doi.org/10.1016/j.fct.2013.01.019>
153. Song X, He Y, Liu M, et al. Mechanism underlying *Polygonum capitatum* effect on *Helicobacter pylori*-associated gastritis based on network pharmacology. *Bioorg Chem.* 2021;114:105044. <https://doi.org/10.1016/j.bioorg.2021.105044>
154. Mendivil EJ, Sandoval-Rodriguez A, Meza-Rios A, et al. Capsaicin induces a protective effect on gastric mucosa along with decreased expression of inflammatory molecules in a gastritis model. *J Funct Foods.* 2019;59:345-351. <https://doi.org/10.1016/j.jff.2019.06.002>
155. Yang HJ, Kim MJ, Kwon DY, et al. Gastroprotective actions of *Taraxacum coreanum* Nakai water extracts in ethanol-induced rat models of acute and chronic gastritis. *J Ethnopharmacol.* 2017;208:84-93. <https://doi.org/10.1016/j.jep.2017.06.045>
156. Jang SW, Lee JW, Park SH, et al. Gastroretentive drug delivery system of DA-6034, a new flavonoid derivative, for the treatment of gastritis. *Int J Pharm.* 2008;356(1-2):88-94. <https://doi.org/10.1016/j.ijpharm.2007.12.042>
157. Chen L, Wei S, He Y, et al. Treatment of chronic gastritis with traditional Chinese medicine: Pharmacological activities and mechanisms. *Pharmaceuticals.* 2023;16:1308. <https://doi.org/10.3390/ph16091308>
158. Carroccio A, Prima LD, Noto D, et al. Searching for wheat plants with low toxicity in celiac disease: Between direct toxicity and immunologic activation. *Digest Liver Dis.* 2011;43(1):34-39. <https://doi.org/10.1016/j.dld.2010.05.005>
159. Karamdoust S, Milani-Hosseini M-R, Faridbod F. Simple detection of gluten in wheat-containing food samples of celiac diets with a novel fluorescent nanosensor made of folic acid-based carbon dots through molecularly imprinted technique. *Food Chem.* 2023;410:135383. <https://doi.org/10.1016/j.foodchem.2022.135383>
160. Birinyi Z, Reder D, Dios A, et al. Immunoanalytic investigation of grain proteins antigenic for celiac disease patients in an einkorn collection. *Food Chem.* 2022;371:131148. <https://doi.org/10.1016/j.foodchem.2021.131148>
161. Dias R, Silva SD, Monteiro B, et al. Mass spectrometry-based quantification of immunostimulatory gliadin proteins and peptides in coloured wheat varieties: Implications for Celiac disease. *Food Res Int.* 2024;178:114008. <https://doi.org/10.1016/j.foodres.2024.114008>
162. Ciacci C, Maiuri L, Caporaso N, et al. Celiac disease: In vitro and in vivo safety and palatability of wheat-free sorghum food products. *Clin Nutr.* 2007;26(6):799-805. <https://doi.org/10.1016/j.clnu.2007.05.006>
163. Mikulska J, Pietrzak D, Rekawek P, et al. Celiac disease and depressive disorders as nutritional implications related to common factors- A comprehensive review. *Behav Brain Res.* 2024;462:114886. <https://doi.org/10.1016/j.bbr.2024.114886>
164. Iacomino G, Aufiero VR, Stasio LD, et al. *Triticum monococcum* amylase trypsin inhibitors possess a reduced potential to elicit innate immune response in celiac patients compared to *Triticum aestivum*. *Food Res Int.* 2021;145:110386. <https://doi.org/10.1016/j.foodres.2021.110386>
165. Malalgoda M, Simsek S. Celiac disease and cereal proteins. *Food Hydrocoll.* 2017;68:108-113. <https://doi.org/10.1016/j.foodhyd.2016.09.024>
166. Pronin D, Borner A, Scherf KA. Old and modern wheat (*Triticum aestivum* L.) cultivars and their potential to elicit celiac disease. *Food Chem.* 2021;339:27952. <https://doi.org/10.1016/j.foodchem.2020.127952>
167. Scherf KA. Immunoreactive cereal proteins in wheat allergy, non-celiac gluten/wheat sensitivity (NCGS) and celiac disease. *Curr Opin Food Sci.* 2019;25:35-41. <https://doi.org/10.1016/j.cofs.2019.02.003>
168. Malalgoda M, Meinhardt SW, Simsek S. Detection and quantitation of immunogenic epitopes related to celiac disease in historical and modern hard red spring wheat cultivars. *Food Chem.* 2018;264:101-107. <https://doi.org/10.1016/j.foodchem.2018.04.131>
169. Victor Z, Julia EH, Tanja S, et al. Viability activation of immune response by quinoa (*Chenopodium quinoa* Willd.) prolamins in celiac disease. *Am J Clin Nutr.* 2012;96(2):337-344. <https://doi.org/10.3945/ajcn.111.030684>
170. Xiao B, Zhang C, Song X, et al. Rationally engineered prolyl endopeptidases from *Sphingomonas capsulata* with improved hydrolytic activity towards pathogenic peptides of celiac disease. *Eur J Med Chem.* 2020;202:112499. <https://doi.org/10.1016/j.ejmech.2020.112499>
171. Maddah R, Panji A, Khoei MA, et al. Deciphering potential biomarkers for celiac disease by using an integrated bioinformatics approach. *Informat Med Unlocked.* 2022;32:101040. <https://doi.org/10.1016/j.imu.2022.101040>

172. Garcia-Calvo E, Garcia-Garcia A, Rodriguez-Gomez S, et al. Development of a new recombinant antibody, selected by phage-display technology from a celiac patient library, for detection of gluten in foods. *Curr Res Food Sci.* 2023;7:100578. <https://doi.org/10.1016/j.crfs.2023.100578>
173. Shoaran M, Sabaie H, Mostafavi M, et al. A comprehensive review of the applications of RNA sequencing in celiac disease research. *Gene.* 2024;927:148681. <https://doi.org/10.1016/j.gene.2024.148681>
174. Singla D, Malik T, Singh A, et al. Advances in understanding wheat-related disorders: A comprehensive review on gluten-free products with emphasis on wheat allergy, celiac and non-celiac gluten sensitivity. *Food Chem Adv.* 2024;4:100627. <https://doi.org/10.1016/j.focha.2024.100627>
175. Sun W, Shahrajabian MH, Cheng Q. Fenugreek cultivation with emphasis on historical aspects and its uses in traditional medicine and modern pharmaceutical science. *Mini Rev Med Chem.* 2021;21(6):724-730. <https://doi.org/10.2174/1389557520666201127104907>
176. Sun W, Shahrajabian MH, Cheng Q. Barberry (*Berberis vulgaris*), a medicinal fruit and food with traditional and modern pharmaceutical uses. *Isr J Plant Sci.* 2021;68(1-2):1-11. <https://doi.org/10.1163/22238980-bja10019>
177. Sun W, Shahrajabian MH, Cheng Q. Health benefits of wolfberry (*Gou Qi Zi*) on the basis of ancient Chinese herbalism and Western modern medicine. *Avicenna J Phytomed.* 2021;11(2):109-119. <https://doi.org/10.22038/AJP.2020.17147>
178. Sun W, Shahrajabian MH, Lin M. Research progress of fermented functional foods and protein factory-microbial fermentation technology. *Fermentation.* 2022;8(12):688. <https://doi.org/10.3390/fermentation8120688>
179. Noori E, Hashemi N, Rezaee D, et al. Potential therapeutic options for celiac disease: An update on current evidence from gluten-free diet to cell therapy. *Int Immunopharmacol.* 2024;133:112020. <https://doi.org/10.1016/j.intimp.2024.112020>
180. Wang N, Cui C, Xu C, et al. Effect and mechanism of peanut skin proanthocyanidins on gliadin-induced Caco-2 celiac disease model cell. *Clin Immunol.* 2022;245:109100. <https://doi.org/10.1016/j.clim.2022.109100>
181. Lavoignat M, Juhasz A, Bose U, et al. Peptidomics analysis of in vitro digested wheat breads: Effect of genotype and environment on protein digestibility and release of celiac disease and wheat allergy related epitopes. *Food Chem.* 2024;448:139148. <https://doi.org/10.1016/j.foodchem.2024.139148>
182. Ribeiro M, Lopes S, Picascia S, et al. Reinventing the nutraceutical value of gluten: The case of L-theanine-gluten as a potential alternative to the gluten exclusion diet in celiac disease. *Food Chem.* 2020;324:126840. <https://doi.org/10.1016/j.foodchem.2020.126840>
183. Schulz K, Giesler L, Linke D, et al. A prolyl endopeptidase from *Flammulina velutipes* for the possible degradation of celiac disease provoking toxic peptides in cereal proteins. *Process Biochem.* 2018;73:47-55. <https://doi.org/10.1016/j.procbio.2018.07.019>
184. Mazzola AM, Zammarchi I, Valerii MC, et al. Gluten-free diet and other celiac disease therapies: Current understanding and emerging strategies. *Nutrients.* 2024;16:1006. <https://doi.org/10.3390/nu16071006>
185. Ahmad AA, Kasim KF, Gopinath SCB, et al. Encapsulation of *Dicranopteris linearis* extract using cellulose microparticles for antiulcer medication. *Int J Biol Macromol.* 2023;253(2):126795. <https://doi.org/10.1016/j.ijbiomac.2023.126795>
186. Nguyen TC, Tang NLC, Le GKN, et al. *Helicobacter pylori* infection and peptic ulcer disease in symptomatic children in Southern Vietnam: A prospective multicenter study. *Healthcare.* 2023;11(11):1658. <https://doi.org/10.3390/healthcare11111658>
187. Noumi E, Dibakto TW. Medicinal plants used for peptic ulcer in the Bangangte region, western Cameroon. *Fitoterapia.* 2000;71(4):406-412. [https://doi.org/10.1016/S0367-326X\(00\)00144-1](https://doi.org/10.1016/S0367-326X(00)00144-1)
188. Yu H-C, Chen T-P, Wei C-Y, et al. Association between peptic ulcer disease and periodontitis: A nationwide population-based case-control study in Taiwan. *Int J Environ Res Public Health.* 2018;15(5):912. <https://doi.org/10.3390/ijreph15050912>
189. Yu Z, Mao X, Tang M, Chen Y, et al. Association between past exposure to fine particulate matter (PM_{2.5}) and peptic ulcer: A cross-sectional study in eastern China. *Chemosphere.* 2021;265:128706. <https://doi.org/10.1016/j.chemosphere.2020.128706>
190. Singh AK, Singh SK, Singh PP, et al. Biotechnological aspects of plants metabolites in the treatment of ulcer: A new prospective. *Biotechnol Rep.* 2018;18:e00256. <https://doi.org/10.1016/j.btre.2018.e00256>
191. Gong H, Zhao N, Zhu C, et al. Treatment of gastric ulcer, traditional Chinese medicine may be a better choice. *J Ethnopharmacol.* 2024;324:117793. <https://doi.org/10.1016/j.jep.2024.117793>
192. Perico LL, Emilio-Silva MT, Ohara R, et al. Systematic analysis of monoterpenes: Advances and challenges in the treatment of peptic ulcer diseases. *Biomolecules.* 2020;10(2):265. <https://doi.org/10.3390/biom10020265>

193. Zhang W, Lian Y, Li Q, Sun L, Chen R, Lai X, Lai Z, Yuan E, Sun S. Preventive and therapeutic potential of flavonoids in peptic ulcers. *Molecules*. 2020;25(20):4626. <https://doi.org/10.3390/molecules25204626>
194. Mahmood S, Fareed MMS, Ahmed G, et al. A robust deep model for classification of peptic ulcer and other digestive tract disorders using endoscopic images. *Biomedicines*. 2022;10(9):2195. <https://doi.org/10.3390/biomedicines10092195>
195. Alomair MK, Alabduladheem LS, Almajed MA, et al. Achillea millefolium essential oil mitigates peptic ulcer in rats through Nrf2/HO-1 pathway. *Molecules*. 2022;27:7908. <https://doi.org/10.3390/molecules2722908>
196. Taha AS, Faccenda E, Angerson WJ, et al. Gastric epithelial anti-microbial peptides-histological correlation and influence of anatomical site and peptic ulcer disease. *Digest Liver Dis*. 2005;37(1):51-56. <https://doi.org/10.1016/j.dld.2004.07.019>
197. Yadav S, Pandey A, Mali SN. From lab to nature: Recent advancements in the journey of gastroprotective agents from medicinal chemistry to phytotherapy. *Eur J Med Chem*. 2024;272:116436. <https://doi.org/10.1016/j.ejmech.2024.116436>
198. Kuna L, Jakab J, Smolic R, et al. Peptic ulcer disease: A brief review of conventional therapy and herbal treatment options. *J Clin Med*. 2019;8(2):179. <https://doi.org/10.3390/jcm8020179>
199. Kumatia EK, Antwi S, Asase A. Analgesic and anti-inflammatory activities of NPK500 capsules, a *Cassia sieberiana* DC.-based herbal analgesic medicine used to treat dysmenorrhea and peptic ulcer, is mediated through the inhibition of PGF2 and iNOS. *J Ethnopharmacol*. 2024. <https://doi.org/10.1016/j.jep.2024.118510>
200. Kumadoh D, Archer M-A, Yeboah GN, et al. A review on anti-peptic ulcer activities of medicinal plants used in the formulation of Enterica, Dyspepsia, and NPK 500 capsules. *Heliyon*. 2021;7:e08465. <https://doi.org/10.1016/j.heliyon.2021.e08465>
201. Huang C-Y, Lai W-Y, Sun M-F, et al. Prescription patterns of traditional Chinese medicine for peptic ulcer disease in Taiwan: A nationwide population-based study. *J Ethnopharmacol*. 2015;176:311-320. <https://doi.org/10.1016/j.jep.2015.11.002>
202. Lakshmi V, Singh N, Shrivastva S, et al. Gedunin and photogedunin of *Xylocarpus granatum* show significant anti-secretory effects and protect the gastric mucosa of peptic ulcer in rats. *Phytomedicine*. 2010;17(8-9):569-574. <https://doi.org/10.1016/j.phymed.2009.10.016>
203. Danai P, Patel S, Pandey V, et al. Antiulcerogenic activity of *Anogeissus pendula* hydroalcoholic extract on pylorus ligated induced gastric ulcers in Albino Wistar rats. *Phytomedicine Plus*. 2021;1(4):100127. <https://doi.org/10.1016/j.phyplu.2021.100127>
204. Jesus NZT, Falcao HS, Filho LMRD, et al. *Hyptis suaveolens* (L.) Poit (Lamiaceae), a medicinal plant protects the stomach against several gastric ulcer models. *J Ethnopharmacol*. 2013;150(3):982-988. <https://doi.org/10.1016/j.jep.2013.10.010>
205. Junior IFDS, Balogun SO, Oliveira RGD, et al. *Piper umbellatum* L.: A medicinal plant with gastric-ulcer protective and ulcer healing effects in experimental rodent models. *J Ethnopharmacol*. 2016;192:123-131. <https://doi.org/10.1016/j.jep.2016.07.011>
206. Simomura VL, Miorando D, Oliveira BMMD, et al. Aqueous extract of the bark of *Uncaria tomentosa*, an Amazonian medicinal plant, promotes gastroprotection and accelerates gastric healing in rats. *J Ethnopharmacol*. 2024;321:117542. <https://doi.org/10.1016/j.jep.2023.117542>
207. Bonamin F, Moraes TM, Santos RCD, et al. The effect of a minor constituent of essential oil from *Citrus aurantium*: The role of β -myrcene in preventing peptic ulcer disease. *Chem Biol Interact*. 2014;212:11-19. <https://doi.org/10.1016/j.cbi.2014.01.009>
208. Memariani Z, Sharifzadeh M, Bozorgi M, et al. Protective effect of essential oil of *Pistacia atlantica* Desf. on peptic ulcer: role of α -pinene. *J Tradit Chin Med*. 2017;37(1):57-63. [https://doi.org/10.1016/S0254-6272\(17\)30027-4](https://doi.org/10.1016/S0254-6272(17)30027-4)
209. Kumar A, Ashwlayan VD, Ratnesh RK, et al. Comparative assessment of anti-ulcer effects: Hydro-ethanolic extract from air-dried ray and disc florets vs essential oil of traditional *Tagetes erecta* (Asteraceae) in Swiss Albino rats. *South Afr J Bot*. 2024;169:197-209. <https://doi.org/10.1016/j.sajb.2024.04.021>
210. Meurer M, Oliveira BMMD, Cury BJ, et al. Extract of *Tagetes erecta* L., a medicinal plant rich in lutein, promotes gastric healing and reduces ulcer recurrence in rodents. *J Ethnopharmacol*. 2022;293:115258. <https://doi.org/10.1016/j.jep.2022.115258>
211. Khan N, Khushhtar M, Rahman MdA, et al. Amelioration of gastric ulcer using a hydro-alcoholic extract of *Mangifera indica* in Sprague Dawley rats by prevention of muco-oxidative stress. *Pharmacol. Res. Modern. Chin Med*. 2024;11:100442. <https://doi.org/10.1016/j.prmcm.2024.100442>

212. Akah PA, Orisakwe OE, Gamaniel KS, et al. Evaluation of Nigerian traditional medicine: II. Effects of some Nigerian folk remedies on peptic ulcer. *J Ethnopharmacol.* 1998;62(2):123-127. [https://doi.org/10.1016/S0378-8741\(98\)00060-9](https://doi.org/10.1016/S0378-8741(98)00060-9)
213. Hamdan DI, Mahmoud MF, Wink M, et al. Effect of hesperidin and neohesperidin from bitter-sweet orange (*Citrus aurantium* var. *bigaradia*) peel on indomethacin-induced peptic ulcers in rats. *Environ. Toxicol Pharmacol.* 2014;37(3):907-915. <https://doi.org/10.1016/j.etap.2014.03.006>
214. Nabil G, Ahmed YH, Ahmed O, et al. Argel, s stemmoside C as a novel natural remedy for mice with alcohol-induced gastric ulcer based on its molecular mechanistic pathway. *J Ethnopharmacol.* 2024;327:117970. <https://doi.org/10.1016/j.jep.2024.117970>
215. Liu H, Chen Y, Hu Y, et al. Protective effects of an alcoholic extract of *Kaempferia galanga* L. rhizome on ethanol-induced gastric ulcer in mice. *J Ethnopharmacol.* 2024;325:117845. <https://doi.org/10.1016/j.jep.2024.117845>
216. Figureiredo FDF, Samazo AS, Arunachalam K, et al. Evaluation of the gastroprotective and ulcer healing properties by *Fridericia chica* (Bonpl.) L.G. Lohmann hydroethanolic extract leaves. *J Ethnopharmacol.* 2023;309:116338. <https://doi.org/10.1016/j.jep.2023.116338>
217. Qu Z, Jiang D, Liu Y, et al. Liuwei Anxiao San protects gastric mucosa from gastric ulcer in rats by regulating the JAK2/STAT3 pathway. *Tissue Cell.* 2023;83:102145. <https://doi.org/10.1016/j.tice.2023.102145>
218. De Jesus NZT, Falcao HDS, Gomes IF, et al. Tannins, peptic ulcers and related mechanisms. *Int J Mol Sci.* 2012;13(3):3203-3228. <https://doi.org/10.3390/ijms13033203>
219. Abdoulrahman K. Anti-ulcer effect of *Ranunculus millefoliatus* on absolute alcohol-induced stomach ulceration. *Saudi J Biol Sci.* 2023;30:103711. <https://doi.org/10.1016/j.sjbs.2023.103711>
220. Ercan G, Tartar RI, Solmaz A, et al. Potent therapeutic effects of ruscogenin on gastric ulcer established by acetic acid. *Asian J Surgery.* 2020;43:405-416. <https://doi.org/10.1016/j.asjsur.2019.07.001>
221. Wu Y-C, Lin L-F, Yeh C-S, et al. Burdock essence promotes gastrointestinal mucosal repair in ulcer patients. *Fooyin J Health Sci.* 2010;2(1):26-31.
222. Yuan E, Lian Y, Li Q, et al. Roles of *Adinandra nitida* (Theaceae) and camellianin A in HCl/ethanol-induced acute gastric ulcer in mice. *Food Sci Human Well.* 2022;11:1053-1063. <https://doi.org/10.1016/j.fshw.2022.03.032>
223. Mohammed DM, Ahmed KA, Desoukey MA, et al. Assessment of the antiulcer properties of *Lawsonia inermis* L. leaves and its nano-formulation against prolonged effect of acute ulcer in rats. *Toxiol Rep.* 2022;9:337-345. <https://doi.org/10.1016/j.toxrep.2022.03.010>
224. Orsi PR, Bonamin F, Severi JA, et al. *Hymenaea stigonocarpa* Mart. ex Hayne: A Brazilian medicinal plant with gastric and duodenal anti-ulcer and antidiarrheal effects in experimental rodent models. *J Ethnopharmacol.* 2012;143:81-90. <https://doi.org/10.1016/j.jep.2012.06.001>
225. Mokam CED, Tamfu AN, Kuissu MM, et al. Phenolic profile and chronic gastric ulcer healing effects of Cameroonian propolis. *Clin Tradit Med Pharmacol.* 2024;5:200140. <https://doi.org/10.1016/j.ctmp.2024.200140>
226. Ofusori AE, Moodley R, Jonnalagadda SB. Antiulcerogenic effects of *Celosia trigyna* plant extracts on ethanol-induced gastric ulcer in adult Wistar rats. *J Tradit Complement Med.* 2020;10:586-593. <https://doi.org/10.1016/j.jtcme.2019.11.004>
227. Algandaby MM. Healing activity of a muco-adhesie formula of *Teucrium polium* against acetic acid-induced oral ulcer in rats. *J King Saud Uni Sci.* 2024;36:103251. <https://doi.org/10.1016/j.jksus.2024.103251>
228. Jabbar AAJ, Mothana RA, Abdulla MA, et al. Mechanisms of anti-ulcer actions of *Prangos pabularia* (L.) in ethanol-induced gastric ulcer in rats. *Saudi Pharm J.* 2023;31:101850. <https://doi.org/10.1016/j.jsps.2023.101850>
229. Hasna B, Houari H, Koula D, et al. In vitro and in vivo study to combined effect of some Algerian medicinal plants and probiotics against *Helicobacter pylori*. *Microorganisms.* 2023;11:1242. <https://doi.org/10.3390/microorganisms11051242>
230. Rosa Junior IA, Almeida DDS, Napolitano HB, et al. Evaluation of gastroprotective activity of the methanolic extract of *Justicia pectoralis* Jacq. (Acanthaceae). *Nutrients.* 2024;16:1430. <https://doi.org/10.3390/nu16101430>
231. Njume C, Jide AA, Ndip RN. Aqueous and organic solvent-extracts of selected South African medicinal plants possess antimicrobial activity against drug-resistant strains of *Helicobacter pylori*: Inhibitory and bactericidal potential. *Int J Mol Sci.* 2011;12:5652-5665. <https://doi.org/10.3390/ijms12095652>
232. Zhang M, Xu L, Chen L, et al. *Dendrobium officinale* polysaccharides as a natural functional component for acetic-acid-induced gastric ulcers in rats. *Molecules.* 2024;29:880. <https://doi.org/10.3390/molecules29040880>

233. Tadesse TY, Zeleke MM, Dagnew SB. Review of ethnobotanical and ethnopharmacological evidence of some Ethiopian medicinal plants traditionally used for peptic ulcer disease treatment. *Clin Experim Gastroenterol.* 2022; 2022:171-187. <https://doi.org/10.2147/CEG.S384395>
234. Sen S, Chakraborty R, De B, et al. Plants and phytochemicals for peptic ulcer: An overview. *Phcog Rev.* 2009;3(6):270-279.
235. Chandrapalan S, Khasawneh F, Singh B, et al. A multi-centre study to risk stratify colorectal polyp surveillance patients utilizing volatile organic compounds and faecal immunochemical test. *Cancers.* 2022;14(19):4951. <https://doi.org/10.3390/cancers14194951>
236. Hsu C-M, Hsu C-C, Hsu Z-M, et al. Colorectal polyp image detection and classification through grayscale images and deep learning. *Sensors.* 2021;21(18):5995. <https://doi.org/10.3390/s21185995>
237. Synytsya A, Vankova A, Miskovicova M, et al. Ex vivo vibration spectroscopic analysis of colorectal polyps for the early diagnosis of colorectal carcinoma. *Diagnostics.* 2021;11(11):2048. <https://doi.org/10.3390/diagnostics11112048>
238. Tse BC, Welham Z, Engel AF, et al. Genomic, microbial and immunological microenvironment of colorectal polyps. *Cancers.* 2021;13(14):3382. <https://doi.org/10.3390/cancers13143382>
239. Yokogawa T, Sasaki Y, Ando H, et al. Pharmacological evaluation for improvement of Kanazawa Sutra, medicinal thread for anal fistula. *J Nat Med.* 2017;71(2):339-348. <https://doi.org/10.1007/s11418-016-1059-0>
240. Mesas C, Quinonero F, Revueltas F, et al. Plant-mediated inorganic nanoparticles for anti-tumor therapy in colorectal cancer: A systematic review. *Appl Sci.* 2023;13(18):10156. <https://doi.org/10.3390/app131810156>
241. Shen M-H, Huang C-C, Chen Y-T, et al. Deep learning empowers endoscopic detection and polyps classification: A multiple-hospital study. *Diagnostics.* 2023;13(8):1473. <https://doi.org/10.3390/diagnostics13081473>
242. Shen N, Ba Q, Lu Y. Causal relationships between polyunsaturated fatty acids and colon polyps: A two-sample mendelian randomization study. *Nutrients.* 2024;16(13):2033. <https://doi.org/10.3390/nu16132033>
243. Abraham A, Jose R, Ahmad J, et al. Comparative analysis of machine learning models for image detection of colonic polyps vs. resected polyps. *J Imaging.* 2023;9(10):215. <https://doi.org/10.3390/jimaging9100215>
244. Dulf E-H, Bleda M, Mocan T, et al. Automatic detection of colorectal polyps using transfer learning. *Sensors.* 2021;21(17):5704. <https://doi.org/10.3390/s21175704>
245. Rejhova A, Opatova A, Cumova A, et al. Natural compounds and combination therapy in colorectal cancer treatment. *Eur J Med Chem.* 2018;144:582-594. <https://doi.org/10.1016/j.ejmech.2017.12.039>
246. Sheikh BY, Sarker MdMR, Kamarudin MNA, et al. Antiproliferative and apoptosis inducing effects of citral via p53 and ROS-induced mitochondrial-mediated apoptosis in human colorectal HCT116 and HT29 cell lines. *Biomed Pharmacother.* 2017;96:834-846. <https://doi.org/10.1016/j.biopha.2017.10.038>
247. Ma F, Lin Y, Ni Z, et al. Therapeutic effects of natural polyphenols on colorectal adenomas: Focus on preclinical studies. *Oncol Rep.* 2023;49(6):112. <https://doi.org/10.3892/or.2023.8549>
248. Esmeeta A, Adhikary S, Dharshna V, et al. Plant-derived bioactive compounds in colon cancer treatment: An updated review. *Biomed Pharmacother.* 2022;153:113384. <https://doi.org/10.1016/j.biopha.2022.113384>
249. Abdullah NA, Hashim NFMD, Zakuan NM, et al. Thioredoxin system in colorectal cancer: Its role in carcinogenesis, disease progression, and response to treatment. *Life Sci.* 2024;348:122711. <https://doi.org/10.1016/j.lfs.2024.122711>
250. Lv S, Ding Y, Huang J, et al. Genetic prediction of micronutrient levels and the risk of colorectal polyps: A mendelian randomization study. *Clin Nutr.* 2024;43(6):1405-1413. <https://doi.org/10.1016/j.clnu.2024.04.019>
251. Islam MdR, Akash S, Rahman MdM, et al. Colon cancer and colorectal cancer: Prevention and treatment by potential natural products. *Chem Biol Interact.* 2022;368:110170. <https://doi.org/10.1016/j.cbi.2022.110170>
252. Hassani S, Maghsoudi H, Fattahi F, et al. Flavonoids nanostructures promising therapeutic efficiencies in colorectal cancer. *Int J Biol Macromol.* 2023;241:124508. <https://doi.org/10.1016/j.ijbiomac.2023.124508>
253. Katharina N, Xuehong Z, Aedin C, et al. Habitual intake of flavonoid subclasses and risk of colorectal cancer in 2 large prospective cohorts. *Am J Clin Nutr.* 2016;103(1):184-191. <https://doi.org/10.3945/ajcn.115.117507>
254. Kong M-Y, Li L-Y, Lou Y-M, et al. Chinese herbal medicines for prevention and treatment of colorectal cancer: From molecular mechanisms of potential clinical applications. *J Integr Med.* 2020;18(5):369-384. <https://doi.org/10.1016/j.joim.2020.07.005>
255. Yang Y, Zhao M, Kuang Q, et al. A comprehensive review of phytochemicals targeting macrophages for the regulation of colorectal cancer progression. *Phytomedicine.* 2024;128:155451. <https://doi.org/10.1016/j.phymed.2024.155451>

256. Bahrami A, Jafari S, Rafiei P, et al. Dietary intake of polyphenols and risk of colorectal cancer and adenoma-A case control study from Iran. *Complement. Ther Med.* 2019;45:269-274. <https://doi.org/10.1016/j.ctim.2019.04.011>
257. Amir M, Ansari MA, Wahab S, et al. In vitro, molecular docking and in silico/ADMET study of cuminaldehyde against Candida, MDR bacteria and human colorectal and cervical carcinoma. *South Afr J Bot.* 2023;163:497-510. <https://doi.org/10.1016/j.sajb.2023.11.011>
258. Weng W, Goel A. Curcumin and colorectal cancer: An update and current perspective on this natural medicine. *Semin Cancer Biol.* 2022;80:73-86. <https://doi.org/10.1016/j.semcancer.2020.02.011>
259. Villegas C, Perez R, Sterner O, et al. Curcuma as an adjuvant in colorectal cancer treatment. *Life Sci.* 2021;286:120043. <https://doi.org/10.1016/j.lfs.2021.120043>
260. Karthika C, Hari B, Mano V, et al. Curcumin as a great contributor for the treatment and mitigation of colorectal cancer. *Experim Gerontol.* 2021;152:111438. <https://doi.org/10.1016/j.exger.2021.111438>
261. Zhong Y, Krisanapun C, Lee S-H, et al. Molecular targets of apigenin in colorectal cancer cells: Involvement of p21, NAG-1 and p53. *Eur J Cancer.* 2010;46(18):3365-3374. <https://doi.org/10.1016/j.ejca.2010.07.007>
262. Cao F, Xia W, Sai S, et al. Berberine: An inspiring resource for the treatment of colorectal diseases. *Biomed Pharmacother.* 2023;167:115571. <https://doi.org/10.1016/j.biopha.2023.115571>
263. Ansari B, Aschner M, Hussain Y, et al. Suppression of colorectal carcinogenesis by naringin. *Phytomedicine.* 2022;96:153897. <https://doi.org/10.1016/j.phymed.2021.153897>
264. Cai K, Cao X-Y, Chen F, et al. Xianlian Jiedu Decoction alleviates colorectal cancer by regulating metabolic profiles, intestinal microbiota and metabolites. *Phytomedicine.* 2024;128:155385. <https://doi.org/10.1016/j.phymed.2024.155385>
265. Fernandez J, Silvan B, Entrialgo-Cadierno R, et al. Antiproliferative and palliative activity of flavonoids in colorectal cancer. *Biomed Pharmacother.* 2021;143:112241. <https://doi.org/10.1016/j.biopha.2021.112241>
266. Cassotta M, Ciansiosi D, Giuseppe RD, et al. Possible role of nutrition in the prevention of inflammatory bowel disease-related colorectal cancer: A focus on human studies. *Nutrition.* 2023;110:111980. <https://doi.org/10.1016/j.nut.2023.111980>
267. Laka K, Mapheto KBF, Mbita Z. Selective in vitro cytotoxicity effect of *Drimys calcarata* bulb extracts against p53 mutant HT-29 and p53 wild-type Caco-2 colorectal cancer cells through STAT5B regulation. *Toxicol Rep.* 2021;8:1265-1279. <https://doi.org/10.1016/j.toxrep.2021.06.015>
268. Ling Q, Fang J, Zhai C, et al. Berberine induces SOCS1 pathway to reprogram the M1 polarization of macrophages via miR-155-5p in colitis-associated colorectal cancer. *Eur J Pharmacol.* 2023;949:175724. <https://doi.org/10.1016/j.ejphar.2023.175724>
269. Nascimento RDPD, Reguengo LM, Machado APDF, et al. The preventive and therapeutic potential of native Brazilian fruits on colorectal cancer. *Food Biosci.* 2022;46:101539. <https://doi.org/10.1016/j.fbio.2021.101539>
270. Qi L, Zhang Y, Song F, Ding Y. Chinese herbal medicine promote tissue differentiation in colorectal cancer by activating HSD11B2. *Arch Biochem Biophys.* 2020;695:108644. <https://doi.org/10.1016/j.abb.2020.108544>
271. Nascimento RDPD, Rizzato JS, Polezi G, et al. Freeze-dried jaboticaba (*Myrciaria jaboticaba*) peel powder, a rich source of anthocyanins and phenolic acids, mitigates inflammation-driven colorectal cancer in mice. *Food Biosci.* 2023;53:102578. <https://doi.org/10.1016/j.fbio.2023.102578>
272. Khan N, Kumar N, Ballal A, et al. Unveiling antioxidant and anti-cancer potentials of characterized *Annona reticulata* leaf extract in 1,2-dimethylhydrazine-induced colorectal cancer in Wistar rats. *J. Ayurveda Integr Med.* 2021;12(4):579-589. <https://doi.org/10.1016/j.jaim.2021.05.010>
273. Nomura AMY, Wilkens LR, Murphy SP, et al. Association of vegetable, fruit, and grain intakes with colorectal cancer: the multiethnic cohort study. *Am J Clin Nutr.* 2008;88(3):730-737. <https://doi.org/10.1093/ajcn/88.3.730>
274. Nirmala P, Ramanathan M. Effect of kaempferol on lipid peroxidation and antioxidant status in 1,2-dimethyl hydrazine induced colorectal carcinoma in rats. *Eur J Pharmacol.* 2011;654(1):75-79. <https://doi.org/10.1016/j.ejphar.2010.11.034>
275. Vuletic S, Bekic M, Tomic S, et al. Could alder buckthorn (*Frangula alnus* Mill.) be a source of chemotherapeutics effective against hepato-and colorectal carcinoma? An in vitro study. *Mutat. Res/Gen. Toxicol Environ Mutagen.* 2023;892:503706. <https://doi.org/10.1016/j.mrgentox.2023.503706>
276. Fernandez J, Saettone P, Franchini MC, et al. Antitumor bioactivity and gut microbiota modulation of polyhydroxybutyrate (PHB) in a rat animal model for colorectal cancer. *Int J Biol Macromol.* 2022;203:638-649. <https://doi.org/10.1016/j.ijbiomac.2022.01.112>

277. Wang D, Zhou Y, Hua L, et al. The role of the natural compound naringenin in AMPK-mitochondria modulation and colorectal cancer inhibition. *Phytomedicine*. 2024;131:155786.
<https://doi.org/10.1016/j.phymed.2024.155786>
278. Macharia JM, Mwangi RW, Rozmann N, et al. Medicinal plants with anti-colorectal cancer bioactive compounds: Potential game-changers in colorectal cancer management. *Biomed Pharmacother*. 2022;153:113383. <https://doi.org/10.1016/j.biopha.2022.113383>
279. Ballou S, Singh P, Nee J, et al. Prevalence and associated factors of bloating: results from the Rome foundation global epidemiology study. *Gastroenterology*. 2023;165(3):647-655.
<https://doi.org/10.1053/j.gastro.2023.05.049>
280. Hussain F, Omran A, Soliman N. Synthesis and characterization of bentonite-based lightweight ceramic aggregates using coal combustion residue and kerosene bloating agent. *Construct Build Mater*. 2024;425:135916. <https://doi.org/10.1016/j.conbuildmat.2024.135916>
281. Palsos OS, Levy RL, Korff MV, et al. Alternative medicine and home remedy use in irritable bowel syndrome (IBS) is related to bloating and diarrhea. *Gastroenterology*. 2003;124(4):A391. [https://doi.org/10.1016/S0016-5085\(03\)81979-0](https://doi.org/10.1016/S0016-5085(03)81979-0)
282. Saiprasad SM, Ramakrishnan M, Savaiano DA. Lactose maldigesters do not adapt to a two-week daily consumption of milk with A1/A2 β -Casein as determined by continued bloating and flatulence. *Curr Develop Nutr*. 2024;8(2):102363. <https://doi.org/10.1016/j.cdnut.2024.102362>
283. Moshiree B, Drossman D, Shaukat A. AGA clinical practice update on evaluation and management of belching, abdominal bloating, and distention: Expert review. *Gastroenterology*. 2023;165(3):791-800.
<https://doi.org/10.1053/j.gastro.2023.04.039>
284. Gundermann D, Ottinger C, Sharp MH, et al. Assessment of the efficacy of a novel oral supplement on symptoms of bloating among healthy pre-menopausal women. *Curr Develop Nutr*. 2024;8(2):102341.
<https://doi.org/10.1016/j.cdnut.2024.102341>
285. Molinari C, Zanelli C, Guarini G, et al. Bloating mechanism in lightweight aggregates: Effect of processing variables and properties of the vitreous phase. *Construct Build Mater*. 2020;261:119980.
<https://doi.org/10.1016/j.conbuildmat.2020.119980>
286. Lacy BE, Cangemi D, Vazquez-Roque M. Management of chronic abdominal distension and bloating. *Clin Gastroenterol Hepatol*. 2021;19(2):219-231. <https://doi.org/10.1016/j.cgh.2020.03.056>
287. Wie YM, Lee KG, Lee KH. Physicochemical effect of the aeration rate on bloating characterizations of artificial lightweight aggregate. *Construct Build Mater*. 2020;256:119444.
<https://doi.org/10.1016/j.conbuildmat.2020.119444>
288. Ha N-Y, Jeong H, Lee H, et al. Safety and effectiveness of traditional herbal medicine Siho-sogan-san in functional dyspepsia: A systematic review and meta-analysis. *J Ethnopharmacol*. 2023;313:116518.
<https://doi.org/10.1016/j.jep.2023.116518>
289. Mahmoudpour Z, Shokri J, Kamalinejad M, et al. The efficacy of a Persian herbal formulation on functional bloating: A double-blind randomized controlled trial. *J Integr Med*. 2019;17(5):344-350.
<https://doi.org/10.1016/j.joim.2019.05.007>
290. Radziszewska M, Smarkusz-Zarzecka J, Ostrowska L. Nutrition, physical activity and supplementation in irritable bowel syndrome. *Nutrients*. 2023;15:3662. <https://doi.org/10.3390/nu15163662>
291. Pop C, Sporea I, Santos J, et al. Efficacy and safety of a novel therapeutics of natural origin (NTN) in adult patients with lactose intolerance: A multicenter, randomized, crossover, double-blind, placebo-controlled study. *Foods*. 2022;11:2600. <https://doi.org/10.3390/foods11172600>
292. Patch C, Pearce AJ, Cheng M, et al. *Bacillus subtilis* (BG0-4TM) improves self-reported symptoms for constipation, indigestion, and dyspepsia: A phase 1/2A randomized controlled trial. *Nutrient*. 2023;15:4490.
<https://doi.org/10.3390/nu15214490>
293. Bertin L, Zanconato M, Crepaldi M, et al. The role of the FODMAP diet in IBS. *Nutrients*. 2024;16:370.
<https://doi.org/10.3390/nu16030370>
294. Giacosa A, Riva A, Petrangolini G, et al. Beneficial effects on abdominal bloating with an innovative food-grade formulation of *Curcuma longa* and *Boswellia serrata* extracts in subjects with irritable bowel syndrome and small bowel dysbiosis. *Nutrients*. 2022;14:416. <https://doi.org/10.3390/nu14030416>
295. Li F, Wang M, Li X, et al. Inflammatory-miR-301a circuitry drives mTOR and Stat3-dependent PSC activation in chronic pancreatitis and PanIN. *Mol Ther Nucl Acids*. 2022;27:970-982.
<https://doi.org/10.1016/j.omtn.2022.01.011>

296. Yin M, Lin J, Wang Y, et al. Development and validation of a multimodel model in predicting severe acute pancreatitis based on radiomics and deep learning. *Int J Med Inform.* 2024;184:105341. <https://doi.org/10.1016/j.ijmedinf.2024.105341>
297. Belabbes FZ.; Mounsif S, Faquir N, et al. Pancreas divisum causing recurrent pancreatitis in a young patient: A case report. *Radiol Case Rep.* 2023;18(1):3535-3538. <https://doi.org/10.1016/j.radcr.2023.07.028>
298. Du D, Yao L, Zhang R, et al. Protective effects of flavonoids from *Coreopsis tinctoria* Nutt. on experimental acute pancreatitis via Nrf-2/ARE-mediated antioxidant pathways. *J Ethnopharmacol.* 2018;224:261-272. <https://doi.org/10.1016/j.jep.2018.06.003>
299. Routh S, Manickam V. Epigenetic alterations dictating the inflammation: A view through pancreatitis. *Life Sci.* 2024;338:122383. <https://doi.org/10.1016/j.lfs.2023.122383>
300. Ko J, Kimita W, Skudder-Hill L, et al. Dietary carbohydrate intake and insulin traits in individuals after acute pancreatitis: Effect modification by intra-pancreatic fat deposition. *Pancreatology.* 2021;21(2):353-362. <https://doi.org/10.1016/j.pan.2021.01.018>
301. Xiang H, Yu H, Zhou Q, et al. Macrophages: A rising star in immunotherapy for chronic pancreatitis. *Pharmacol Res.* 2022;185:106508. <https://doi.org/10.1016/j.phrs.2022.106508>
302. Zhou Y, Huang X, Jin Y, et al. The role of mitochondrial damage-associated molecular patterns in acute pancreatitis. *Biomed Pharmacother.* 2024;175:116690. <https://doi.org/10.1016/j.biopha.2024.116690>
303. Huang W, Zhang J, Jin W, et al. Piperine alleviates acute pancreatitis: A possible role for FAM134B and CCPG1 dependent ER-phagy. *Phytomedicine.* 2022;105:154361. <https://doi.org/10.1016/j.phymed.2022.154361>
304. Yuan C, Dong X, Xu S, et al. AKBA alleviates experimental pancreatitis by inhibiting oxidative stress in macrophages through the Nrf2/HO-1 pathway. *Int Immunopharmacol.* 2023;121:110501. <https://doi.org/10.1016/j.intimp.2023.110501>
305. Ramakrishnan P, Loh WM, Gopinath SCB, et al. Selective phytochemicals targeting pancreatic stellate cells as new anti-fibrotic agents for chronic pancreatitis and pancreatic cancer. *Acta Pharm Sin B.* 2020;10(3):399-413. <https://doi.org/10.1016/j.apsb.2019.11.008>
306. Tang Y, Sun M, Liu Z. Phytochemicals with protective effects against acute pancreatitis: a review of recent literature. *Pharmaceut Biol.* 2022;60(1):479-490. <https://doi.org/10.1080/13880209.2022.2039723>
307. Choi J-W, Shin J, Zhou Z, et al. Myricetin ameliorates the severity of pancreatitis in mice by regulating cathepsin B activity and inflammatory cytokine production. *Int Immunopharmacol.* 2024;136:112284. <https://doi.org/10.1016/j.intimp.2024.112284>
308. Zhou L, Yu J, Wang S, et al. Tectoridin alleviates caerulein-induced severe acute pancreatitis by targeting ERK2 to promote macrophage M2 polarization. *Arch Biochem Biophys.* 2024;752:109873. <https://doi.org/10.1016/j.abb.2023.109873>
309. Chelpuri Y, Pabbathi S, Alla GR, et al. Tropolone derivative hinokitiol ameliorates cerulein-induced acute pancreatitis in mice. *Int Immunopharmacol.* 2022;109:108915. <https://doi.org/10.1016/j.intimp.2022.108915>
310. Han L, Faap MD. Original research: *Bacillus coagulans* significantly improved abdominal pain and bloating in patients with IBS. *Postgraduate Med.* 2009;121(2):119-124. <https://doi.org/10.3810/pgm.2009.03.1984>
311. Huang Y, Wen Y, Wang R, et al. Temporal metabolic trajectory analyzed by LC-MS/MS based targeted metabolomics in acute pancreatitis pathogenesis and Chaiqin Chengqi decoction therapy. *Phytomedicine.* 2022;99:153996. <https://doi.org/10.1016/j.phymed.2022.153996>
312. Cao F, Xiang J, Wang Y, et al. Chaiqin chengqi decoction alleviates acute pancreatitis by targeting gasdermin D-mediated pyroptosis. *J Ethnopharmacol.* 2024;318:116920. <https://doi.org/10.1016/j.jep.2023.116920>
313. Wen Y, Han C, Liu T, et al. Chaiqin chengqi decoction alleviates severity of acute pancreatitis vis inhibition of TLR4 and NLRP3 inflammasome: Identification of bioactive ingredients via pharmacological sub-network analysis and experimental validation. *Phytomedicine.* 2020;79:153328. <https://doi.org/10.1016/j.phymed.2020.153328>
314. Han X, Bao J, Ni J, et al. Qing Xia Jie Yi formular granules alleviated acute pancreatitis through inhibition of M1 macrophage polarization by suppressing glycolysis. *J Ethnopharmacol.* 2024;325:117750. <https://doi.org/10.1016/j.jep.2024.117750>
315. Huang Y-T, Liang Q-Q, Zhang H-R, et al. Baicalin inhibits necroptosis by decreasing oligomerization of phosphorylated MLKL and mitigates caerulein-induced acute pancreatitis in mice. *Int Immunopharmacol.* 2022;108:108885. <https://doi.org/10.1016/j.intimp.2022.108885>

316. Abozaid OAR, Moawed FSM, Ahmed ESA, et al. Cinnamic acid nanoparticles modulate redox signal and inflammatory response in gamma irradiated rats suffering from acute pancreatitis. *Biochim. Biophys. Acta Mol. Basis Dis.* 2020; 1866(11), 165904. <https://doi.org/10.1016/j.bbadis.2020.165904>
317. Bansod S, Doijad N, Godugu C. Berberine attenuates severity of chronic pancreatitis and fibrosis via AMPK-mediated inhibition of TGF- β 1/Smad signaling and M2 polarization. *Toxicol Appl Pharmacol.* 2020;403:115162. <https://doi.org/10.1016/j.taap.2020.115162>
318. Aja PM, Izekwe FI, Famurewa AC, et al. Hesperidin protects against cadmium-induced pancreatitis by modulating insulin secretion, redox imbalance, and iNOS/NF- κ B signaling in rats. *Life Sci.* 2020;259:118268. <https://doi.org/10.1016/j.lfs.2020.118268>
319. Fan R, Sui J, Dong X, et al. Wedelolactone alleviates acute pancreatitis and associated lung injury via GPX4 mediated suppression of pyroptosis and ferroptosis. *Free Radic Biol Med.* 2021;173:29-40. <https://doi.org/10.1016/j.freeradbiomed.2021.07.009>
320. Lin Y-P, Ni Y-H, Weng W-C, et al. Cyclic vomiting syndrome and migraine in children. *J Formosan Med Associat.* 2011;110(6):382-387. [https://doi.org/10.1016/S0929-6646\(11\)60056-9](https://doi.org/10.1016/S0929-6646(11)60056-9)
321. Jin H, Zhao K, Li J, et al. Matrine alleviates oxidative stress and ferroptosis in severe acute pancreatitis-induced acute lung injury by activating the UCP2/SIRT3/PGC1 α pathway. *Int Immunopharmacol.* 2023;117:109981. <https://doi.org/10.1016/j.intimp.2023.109981>
322. Ohkawara T, Takeda H, Nishihira J. Protective effect of chlorogenic acid on the inflammatory damage of pancreas and lung in mice with L-arginine-induced pancreatitis. *Life Sci.* 2017;190:91-96. <https://doi.org/10.1016/j.lfs.2017.09.015>
323. Liang X, Hu C, Liu C, et al. Dihydrokaempferol (DHK) ameliorates severe acute pancreatitis (SAP) via Keap1/Nrf2 pathway. *Life Sci.* 2020;261:118340. <https://doi.org/10.1016/j.lfs.2020.118340>
324. Chen Y-J, Chen M-Z, Zhang H-W, et al. Effect of Guo Qing Yi Tang combined with Western medicine cluster therapy on acute pancreatitis. *Am J Emerg Med.* 2021;50:66-70. <https://doi.org/10.1016/j.ajem.2021.07.004>
325. Silva-Leite KESD, Girao DKFB, Pires ADF, et al. Ximenia americana heteropolysaccharides ameliorate inflammation and visceral hypernociception in murine caerulein-induced acute pancreatitis: Involvement of CB2 receptors. *Biomed Pharmacother.* 2018;106:1317-1324. <https://doi.org/10.1016/j.biopha.2018.07.067>
326. Yuvaraj D, Arumugam G. Anti-inflammatory and preventive activity of white mulberry root bark extract in an experimental model of pancreatitis. *J Tradit Complement Med.* 2018;8:497-505. <https://doi.org/10.1016/j.jtcme.2018.01.011>
327. Lima PR, Melo TSD, Carvalho KMMB, et al. 1,8-cineole (eucalyptol) ameliorates cerulein-induced acute pancreatitis via modulation of cytokines, oxidative stress and NF- κ B activity in mice. *Life Sci.* 2013;92:1195-1201. <https://doi.org/10.1016/j.lfs.2023.05.009>
328. Zhang J, Wang Y, Liu B, et al. Activity of Ligustrum vulgare L. extract against acute pancreatitis in murine models by regulation of p38 MAPK and NF- κ B signaling pathways. *Saudi J Biol Sci.* 2022;29:273-278. <https://doi.org/10.1016/j.sjbs.2021.08.087>
329. Suguna P, Geetha A, Aruna R, et al. Nigella sativa Linn. seed extract modulates the activity of ASC complex of NLRP3 inflammasome in rats subjected to experimental pancreatitis. *Biomed Prevent Nutr.* 2014;4(2):113-120. <https://doi.org/10.1016/j.bionut.2013.12.008>
330. Lampropoulos P, Lambropoulou M, Papalois A, et al. The role of apigenin in an experimental model of acute pancreatitis. *J. Surg. Res.* 2013;183(1):129-137. <https://doi.org/10.1016/j.jss.2012.11.053>
331. Seo JY, Pandey RP, Lee J, et al. Quercetin 3-O-xyloside ameliorates acute pancreatitis in vitro via the reduction of ER stress and enhancement of apoptosis. *Phytomedicine.* 2019;55:40-49. <https://doi.org/10.1016/j.phymed.2018.07.011>
332. Bae G-S, Kim M-S, Jeong J, et al. Piperine ameliorates the severity of cerulein-induced acute pancreatitis by inhibiting the activation of mitogen activated protein kinases. *Biochem Biophys Res Commun.* 2011;410(3):382-388 <https://doi.org/10.1016/j.bbrc.2011.05.136>
333. Jo I-J, Bae G-S, Choi SB, et al. Fisetin attenuates cerulein-induced acute pancreatitis through down regulation of JNK and NF- κ B signaling pathways. *Eur J Pharmacol.* 2014;737:149-158. <https://doi.org/10.1016/j.ejphar.2014.05.018>
334. Zhao X-L, Xiang J, Wan M-H, et al. Effect of acute pancreatitis on the pharmacokinetics of Chinese herbal ointment Liu-He-Dan in anaesthetized rats. *J Ethnopharmacol.* 2013;145(1):94-99. <https://doi.org/10.1016/j.jep.2012.10.036>

335. Anchi P, Khurana A, Swain D, et al. Sustained-release curcumin microparticles for effective prophylactic treatment of exocrine dysfunction of pancreas: A preclinical study on cerulein-induced acute pancreatitis. *J Pharma Sci.* 2018;107(11):2869-2882. <https://doi.org/10.1016/j.xphs.2018.07.009>
336. Choi S-B, Bae G-S, Jo I-J, et al. Berberine inhibits inflammatory mediators and attenuates acute pancreatitis through deactivation of JNK signaling pathways. *Mol Immunol.* 2016;74:27-38. <https://doi.org/10.1016/j.molimm.2016.04.011>
337. Xia Y, Xu Y, Liu Q, et al. Glutaredoxin 1 regulates cholesterol metabolism and gallstone formation by influencing protein S-glutathionylation. *Metabolism.* 2023;145:155610. <https://doi.org/10.1016/j.metabol.2023.155610>
338. Kim D-G, Bae G-S, Choi S-B, et al. Guggulsterone attenuates cerulein-induced acute pancreatitis via inhibition of ERK and JNK activation. *Int Immunopharmacol.* 2015;26(1):194-202. <https://doi.org/10.1016/j.intimp.2015.03.030>
339. Bansod S, Godugu C. Nimbolide ameliorates pancreatic inflammation and apoptosis by modulating NF- κ B/SIRT1 and apoptosis signaling in acute pancreatitis model. *Int Immunopharmacol.* 2021;90:107264. <https://doi.org/10.1016/j.intimp.2020.107264>
340. Kouchi K, Takenouchi A, Matsuoka A, et al. Efficacy of an anal fistula plug for fistulas-in-Ano in children. *J Pediatric Surgery.* 2017;52(8):1280-1282. <https://doi.org/10.1016/j.jpedsurg.2017.01.062>
341. Du D, Wang X, Zuo H, et al. Differences in glucose homeostasis and islet injury among diverse mice strains post acute pancreatitis. *Biochem Biophys Res Commun.* 2024;708:149780. <https://doi.org/10.1016/j.bbrc.2024.149780>
342. Das D, Mukherjee S, Das AS, et al. Aqueous extract of black tea (*Camellia sinensis*) prevents ethanol+cholecystokinin-induced pancreatitis in a rat model. *Life Sci.* 2006;78(19):2194-2203. <https://doi.org/10.1016/j.lfs.2005.09.020>
343. Bashir N, Manoharan V, Miltonprabu S. Grape seed proanthocyanidins protects against cadmium induced oxidative pancreatitis in rats by attenuating oxidative stress, inflammation and apoptosis via Nrf-2/HO-1 signaling. *J Nutr Biochem.* 2016;32:128-141. <https://doi.org/10.1016/j.jnutbio.2016.03.001>
344. Santana DG, Santos CA, Santos ADC, et al. Beneficial effects of the ethanol extract of *Caesalpinia pyramidalis* on the inflammatory response and abdominal hyperalgesia in rats with acute pancreatitis. *J Ethnopharmacol.* 2012;142:445-455.
345. Pasari LP, Khurana A, Anchi P, et al. Visnagin attenuates acute pancreatitis via Nrf2/NF κ B pathway and abrogates associated multiple organ dysfunction. *Biomed Pharmacother.* 2019;112:108629. <https://doi.org/10.1016/j.biopha.2019.108629>
346. Setia G, Ramanathan M, Bhavanani AB, et al. Adjuvant yoga therapy for symptom management of functional dyspepsia: A case series. *J Ayurveda Integr Med.* 2023;14(3):100715. <https://doi.org/10.1016/j.jaim.2023.100715>
347. Boyd T, Paz M, Ahmad I, et al. Unrecognized functional dyspepsia among those with refractory chronic constipation: Analysis of a tertiary cohort. *Gastro Hep Adv.* 2023;2(4):573-579. <https://doi.org/10.1016/j.gastha.2023.01.004>
348. Correia H, Peneiras S, Levchuk N, et al. Effects of a non-caffeinated coffee substitute on functional dyspepsia. *Clin Nutr ESPEN.* 2021;41:412-416. <https://doi.org/10.1016/j.clnesp.2020.10.009>
349. Guo Y, Zhu J, Su X, et al. Efficacy of Chinese herbal medicine in functional dyspepsia: A meta-analysis of randomized, double-blind, placebo-controlled trials. *J Tradit Chin Med Sci.* 2016;3:147-156. <https://doi.org/10.1016/j.jtcms.2016.09.006>
350. Alexander CF, Mahadeva S, Carbone F, et al. Functional dyspepsia. *The Lancet.* 2020;396(10263):1689-1702. [https://doi.org/10.1016/S0140-6736\(20\)30469-4](https://doi.org/10.1016/S0140-6736(20)30469-4)
351. Ullah H, Minno AD, Piccinocchi R, et al. Efficacy of digestive enzyme supplementation in functional dyspepsia: A monocentric, randomized, double-blind, placebo-controlled, clinical trial. *Biomed Pharmacother.* 2023;169:115858. <https://doi.org/10.1016/j.biopha.2023.115858>
352. Chen S-H, Wu H-S, Jiang X-F, et al. Bioinformatics and LC-QTOF-MS based discovery of pharmacodynamic and Q-markers of Pitongshu against functional dyspepsia. *J Ethnopharmacol.* 2024;329:118096. <https://doi.org/10.1016/j.jep.2024.118096>
353. Ceulemans M, Wauters L, Vanuytsel T. Targeting the altered duodenal microenvironment in functional dyspepsia. *Curr Opin Pharmacol.* 2023;70:102363. <https://doi.org/10.1016/j.coph.2023.102363>
354. Sato H, Grover M. Gastroparesis and functional dyspepsia: Spectrum of gastroduodenal neuromuscular disorders or unique entities? *Gastro Hep Adv.* 2023;2(3):438-448. <https://doi.org/10.1016/j.gastha.2022.10.005>

355. Zheng Y-F, Liang S-P, Zhong Z-S, et al. Duodenal microbiota makes an important impact in functional dyspepsia. *Microb Pathogen*. 2022;162:105297. <https://doi.org/10.1016/j.micpath.2021.105297>
356. Murray HB, Ljotsson B, Healy B, et al. Brief cognitive-behavioral treatment for avoidant/restrictive food intake disorder in the context of functional dyspepsia: Study protocol for a feasibility randomized controlled trial. *Centemporary Clin Trials*. 2023;134:107336. <https://doi.org/10.1016/j.cct.2023.107336>
357. Ho LTF, Chung VCH, Wong CHL, et al. Evaluating traditional Chinese medicine diagnostic instruments for functional dyspepsia: A systematic review on measurement properties. *Integr Med Res*. 2021;10:100713. <https://doi.org/10.1016/j.jmr.2020.100713>
358. Luo X, Yang Y, Mao X, et al. Traditional Chinese medicine compounds for the treatment of functional dyspepsia: An updated meta-analysis of randomized, double-blind, placebo-controlled trials. *Digital Chin Med*. 2021;4:273-289.
359. Popa SL, Dumitrascu DI, Pop C, et al. Exclusion diets in functional dyspepsia. *Nutrients*. 2022;14:2057. <https://doi.org/10.3390/nu14102057>
360. Liao K-Y, Chang F-Y, Wu L-T, et al. Cyclic vomiting syndrome in Taiwanese children. *J Formosan Med Associat*. 2011;110(1):14-18. [https://doi.org/10.1016/S0929-6646\(11\)60003-X](https://doi.org/10.1016/S0929-6646(11)60003-X)
361. Zhang J, Chen T, Wen Y, et al. Insights and future prospects of traditional Chinese medicine in the treatment of functional dyspepsia. *Phytomedicine*. 2024;127:155481. <https://doi.org/10.1016/j.phymed.2024.155481>
362. Zeng Y, Zhou L, Wan Y, et al. Effect of Saikosaponin D on apoptosis, autophagy, and morphological structure of intestinal cells of Cajal with functional dyspepsia. *Combinat. Chem. High. Throughput Screen*. 2024;27(10):1513-1522. <https://doi.org/10.2174/0113862073262404231004053116>
363. Liu T, Asif IM, Liu L, et al. Laminarin ameliorates iodacetamide-induced functional dyspepsia via modulation of 5-HT₃ receptors and the gut microbiota. *Int J Biol Macromol*. 2024;268(1):131640. <https://doi.org/10.1016/j.ijbiomac.2024.131640>
364. Wang Y, Yao J, Zhu Y, et al. Combination of Simo Decoction and Golden Bifid alleviates functional dyspepsia through a mechanism involving intestinal microbiota and short-chain fatty acids. *Arab J Gastroenterol*. 2024. <https://doi.org/10.1016/j.ajg.2023.12.009>
365. Liu R, Li T, Xu H, et al. Systems biology strategy through integrating metabolomics and network pharmacology to reveal the mechanisms of Xiaopi Hewei Capsule improves functional dyspepsia. *J Chromatogr B*. 2023;1226:123676. <https://doi.org/10.1016/j.jchromb.2023.123676>
366. Bai M, Zhao L, Liu M, et al. Deciphering the function of Xiangsha-Liujunzi-Tang in enhancing duodenal mucosal barrier by inhibiting MC/Tryptase/PAR-2 signaling pathway in functional dyspepsia rats. *J Ethnopharmacol*. 2024;319:116715. <https://doi.org/10.1016/j.jep.2023.116715>
367. Sun J, Gou J, Qin L, et al. Screening of anti-functional dyspepsia compounds in *Cynanchum auriculatum*: A spectrum-effect relationship analysis, and ATP-binding cassette transporters inhibitor evaluation. *J Ethnopharmacol*. 2024;318:116867. <https://doi.org/10.1016/j.jep.2023.116867>
368. Zhang D, Gu J, Xu Y, et al. Exploring the mechanism of Huanglian ointment in alleviating wound healing after anal fistula surgery through metabolomics and proteomics. *Heliyon*. 2024;10:e29809. <https://doi.org/10.1016/j.heliyon.2024.e29809>
369. Moghtaderi S, Rameshk M, Hasheminasab FS, et al. Distribution of traditional Persian medicine stomach dysmperaments in patients with functional dyspepsia: A cross-sectional study. *Eur J Integr Med*. 2023;58:102215. <https://doi.org/10.1016/j.eujim.2022.102215>
370. Ho L, Zhong CCW, Wong CHL, et al. Herbal medicine for functional dyspepsia: Network meta-analysis of placebo-controlled randomized trials. *J Ethnopharmacol*. 2022;283:114665. <https://doi.org/10.1016/j.jep.2021.114665>
371. Gan Y, Liu H, Yang L, et al. Effect of Banxiaxixin Tang on treatment of functional dyspepsia: a Meta-analysis of randomized controlled trials. *J Tradit Chin Med*. 2014;34(2):140-144.
372. Mohtashami R, Huseini HF, Heydari M, et al. Efficacy and safety of honey based formulation of *Nigella sativa* seed oil in functional dyspepsia: A double blind randomized controlled clinical trial. *J Ethnopharmacol*. 2015;175:147-152. <https://doi.org/10.1016/j.jep.2015.09.022>
373. Khonche A, Huseini HF, Abdi H, et al. Efficacy of *Mentha pulegium* extract in the treatment of functional dyspepsia: A randomized double-blind placebo-controlled clinical trial. *J Ethnopharmacol*. 2017;206:267-273. <https://doi.org/10.1016/j.jep.2017.05.026>

374. Tu Y, Luo X, Liu D, et al. Extracts of *Poria cocos* improve functional dyspepsia via regulating brain-gut peptides, immunity and repairing of gastrointestinal mucosa. *Phytomedicine*. 2022;95:153875. <https://doi.org/10.1016/j.phymed.2021.153875>
375. Rosch W, Liebrechts T, Gundermann K-J, et al. Phytotherapy for functional dyspepsia: A review of the clinical evidence for the herbal preparation STW5. *Phytomedicine*. 2006;13(1):114-121. <https://doi.org/10.1016/j.phymed.2006.03.002>
376. Gao S, Huang Q, Tang X, et al. ZhiShiXiaoPi tang inhibits autophagy induced by corticosterone and functional dyspepsia through blockade of the mTOR pathway. *J Ethnopharmacol*. 2019;238:111836. <https://doi.org/10.1016/j.jep.2019.111836>
377. Mobeen A, Hakeem A. Efficacy of herbal formulations Sufoof-i-Tabkheer and Arq-i-Badiyan in dyspepsia: An open-labeled single-arm clinical study. *J Herb Med*. 2023;41:100700. <https://doi.org/10.1016/j.hermed.2023.100700>
378. Wang Y, Jia Y, Liu X, et al. Effect of Chaihu-Shugan-San on functional dyspepsia and gut microbiota: A randomized, double-blind, placebo-controlled trial. *J Ethnopharmacol*. 2024;322:117659. <https://doi.org/10.1016/j.jep.2023.117659>
379. Jin C, Zhang Z, Li X. Silver nanoparticles stabilized by chitosan-polyvinyl alcohol polymers mediated by Pistacia extract for treatment of functional dyspepsia and gastric cancer. *J Sci Adv Mater Device*. 2024;9:100757. <https://doi.org/10.1016/j.jsamd.2024.100757>
380. Qiankun L, Lanfang M, Xiaojuan D, et al. Pingwei capsules improve gastrointestinal motility in rats with functional dyspepsia. *J Tradit Chin Med*. 2018;38(1):43-53.
381. Li X, Lin Y, Jiang Y, et al. Aqueous extract of *Phyllanthus emblica* L. alleviates functional dyspepsia through regulating gastrointestinal hormones and gut microbiota in vivo. *Foods*. 2022;11:1491. <https://doi.org/10.3390/foods11101491>
382. Choi N-R, Jung D, Kim S-C, et al. Analysis of network pharmacological efficacy and therapeutic effectiveness in animal models for functional dyspepsia of *Foeniculi fructus*. *Nutrients*. 2023;15:2644. <https://doi.org/10.3390/nut15122644>
383. Lee B, Ha N-Y, Park H-J, et al. Herbal medicine Yukgunja-Tang for functional dyspepsia: A protocol for a randomized, controlled, multicenter clinical trial. *Healthcare*. 2023;11:1456. <https://doi.org/10.3390/healthcare11101456>
384. Zhang J, Wang X, Wang F, et al. Xiangsha Liujunzi decoction improves gastrointestinal motility in functional dyspepsia with spleen deficiency syndrome by restoring mitochondrial quality control homeostasis. *Phytomedicine*. 2022;105:154374. <https://doi.org/10.1016/j.phymed.2022.154374>
385. Li Z, Yang T, Wang L, et al. Comparison of the effects of *Amomum tsaoko* and its adulterants on functional dyspepsia rats based on metabolomics analysis. *J Pharm Biomed Anal*. 2024;246:116208. <https://doi.org/10.1016/j.jpba.2024.116208>
386. Liu B, Kou Z, Chen B. Effects and mechanisms of traditional Chinese medicines on functional dyspepsia: A review. *Chin Herb Med*. 2023;15:516-525. <https://doi.org/10.1016/j.chmed.2023.06.001>
387. Lin J, Han X, Dai N, et al. Effectiveness of Chengqi-series decoctions in treating severe acute pancreatitis: A systematic review and meta-analysis. *Phytomedicine*. 2023;113:154727. <https://doi.org/10.1016/j.phymed.2023.154727>
388. Rafieian-kopaei M, Hosseini-asl K. Effects of *Ocimum basilicum* on functional dyspepsia: a double-blind placebo-controlled study. *Iran J Med Sci*. 2005;30(3):134-137.
389. Gallo A, Pellegrino S, Lipari A, et al. Lactose malabsorption and intolerance: What is the correct management in older adults? *Clin Nutr*. 2023;42(12):2540-2545. <https://doi.org/10.1016/j.clnu.2023.10.014>
390. Wang J, Sijs B, Bakker W, et al. Ribotoxin deoxynivalenol induces taurocholic acid malabsorption in an in vitro human intestinal model. *Toxicol Lett*. 2023;383:54-63. <https://doi.org/10.1016/j.toxlet.2023.06.001>
391. Mergener R, Nunes MR, Nascimento LPC, et al. Congenital glucose-galactose malabsorption: A case report about cause and consequence, not exactly in this order. *Global Pediatrics*. 2024;9:100181. <https://doi.org/10.1016/j.gpeds.2024.100181>
392. Medernach J, Middleton JP. Malabsorption syndromes and food intolerance. *Clin Perinatol*. 2022;49(2):537-555. <https://doi.org/10.1016/j.clp.2022.02.015>
393. Born P. The clinical impact of carbohydrate malabsorption. *Arab J Gastroenterol*. 2011;12(1):1-14. <https://doi.org/10.1016/j.ajg.2011.01.002>

394. Evenson A, Thompson O, Crema A. The relationship of anxiety and stress to lactose malabsorption and gastrointestinal symptoms. *J Acad Nutr. Diet.* 2023;123(9):A40. <https://doi.org/10.1016/j.jand.2023.06.135>
395. Montoro-Huguet MA, Belloc B, Dominguez-Cajal M. Small and large intestine (I): Malabsorption of nutrients. *Nutrients.* 2021;13(4):1254. <https://doi.org/10.3390/nu13041254>
396. Seth A, Padda I, Chaudhry R. A rare complication of secondary pneumothorax following exploratory laparotomy for a small bowel obstruction: Case report. *Int J Surgery Case Rep.* 2023;102:107825. <https://doi.org/10.1016/j.ijscr.2022.107825>
397. Khanal N, Thapa A, Raut A, et al. Intra-abdominal propylene structure fragment leading to complete small bowel obstruction: A case report. *Int J Surgery Case Rep.* 2021;85:106174. <https://doi.org/10.1016/j.ijscr.2021.106174>
398. Bauschert L, Sermet K, Frealle E, et al. A case of subacute bowel obstruction revealing slowly-evolutive gastro-intestinal mucormycosis following allogeneic hematopoietic cell transplantation. *J Med Mycol.* 2022;32(4):101312. <https://doi.org/10.1016/j.mycmed.2022.101312>
399. Peters PN, Moyett JM, Davidson BA, et al. Cost-effectiveness of management strategies for patients with recurrent ovarian cancer and inoperable malignant bowel obstruction. *Gynecol Oncol.* 2022;167(3):523-531. <https://doi.org/10.1016/j.ygyno.2022.10.013>
400. Shrestha BM, Shrestha S, Aryal S, et al. Transmesenteric hernia with closed-loop small bowel obstruction: A case report. *Annal Med Surgery.* 2022;74:103256. <https://doi.org/10.1016/j.amsu.2022.103256>
401. Knight HP, Hollahan CG, Leung SOA, et al. Implementing a clinical pathway for malignant bowel obstruction in patients with gynecologic malignancies. *J Pain Symptom Manag.* 2024;67(5):e741. <https://doi.org/10.1016/j.jpainsymman.2024.02.234>
402. Pakhare P, Milan, Ghimire S, et al. Complete small bowel obstruction due to ileal volvulus with autosomal dominant polycystic kidney disease: A rare case report. *Int J Surgery Case Rep.* 2023;105:108052. <https://doi.org/10.1016/j.ijscr.2023.108052>
403. Zhang S, Chen H-Y, Shao B, et al. Effect of herbs on postoperative intestinal obstruction: A case report. *Altern Ther Health Med.* 2021;27(5):74-76.
404. Kunitomi Y, Nakashima M, Takeuchi M, et al. Efficacy of Daikenchuto in the prevention of bowel obstruction in patients with colorectal cancer undergoing laparoscopic surgery: An observational study using a Japanese administrative claims database. *Support Care Cancer.* 2023;31(2):133. <https://doi.org/10.1007/s00520-023-07599-5>
405. Yu X, Cui Z, Zhou Z, et al. Si-Jun-Zi decoction treatment promotes the restoration of intestinal function after obstruction by regulating intestinal homeostasis. *Evid Based Complement Altern Med.* 2014;2014(11):928579. <https://doi.org/10.1155/2014/928579>
406. Emile SH, Magdy A, Youssef M, et al. Utility of endoanal ultrasonography in assessment of primary and recurrent anal fistulas and for detection of associated anal sphincter defects. *J Gastrointestinal Surgery.* 2017;21(11):1879-1887. <https://doi.org/10.1007/s11605-017-3574-z>
407. Kumar MM, Tamizhselvi R. Protective effect of diallyl disulfide against cerulein-induced acute pancreatitis and associated lung injury in mice. *Int Immunopharmacol.* 2020;80:106136. <https://doi.org/10.1016/j.intimp.2019.106136>
408. Mendes CRS, Ferreira LSDC, Sapucaia RA, et al. VAAFT- Videoassisted anal fistula treatment: A new approach for anal fistula VAAFT- Tratamento de fistula anal videoassistida: uma nova abordagem para a fistula anal. *J Coloproctol.* 2014;34(1):62-64. <https://doi.org/10.1016/j.jcol.2014.02.005>
409. Garg P.; Sachdeva M, Yagnik VD. TROPIS (Transanal Opening of Intersphincteric Space) procedure for the treatment of horseshoe anal fistulas. *J Gastrointestinal Surgery.* 2022;26(8):1814-1815. <https://doi.org/10.1007/s11605-022-05384-z>
410. Abdelnaby M, Emile S, El-Said M, et al. Drained mucosal advancement flap versus rerouting Seton around the internal anal sphincter in treatment of high trans-sphincteric anal fistula: A randomized trial. *Int J Surgery.* 2019;72:198-203. <https://doi.org/10.1016/j.ijssu.2019.11.008>
411. Wright M, Thorson A, Blatchford G, et al. What happens after a failed LIFT for anal fistula? *Am J Surgery.* 2017;214(6):1210-1213. <https://doi.org/10.1016/j.amjsurg.2017.08.042>
412. Lynn PB, Carrano FM, Grieco M, et al. Ligation of the intersphincteric Fistula tract (LIFT) as a first approach in the surgical treatment of transphincteric anal fistula is associated with modest initial success rates. *Surgery Open Digest Adv.* 2023;9:100077. <https://doi.org/10.1016/j.soda.2022.100077>

413. Pigot F. Treatment of anal fistula and abscess. *J Visceral Surgery*. 2015;152(2):S23-S29.
<https://doi.org/10.1016/j.jviscsurg.2014.07.008>
414. Kumar A, Sharma A, Gupta SJ, et al. Complex anal fistula treated with IFTAK (Interception of fistulous track with application of ksharsutra) technique- A case report. *J Ayurveda Integr Med*. 2023;14(1):100686.
<https://doi.org/10.1016/j.jaim.2023.100686>
415. Ellis CN. Bioprosthetic plugs for complex anal fistulas: An early experience. *J Surgical Educat*. 2007;64(1):36-40.
<https://doi.org/10.1016/j.cursur.2006.07.005>
416. Gou B, Zhang J-C, Chen L, et al. Comparison of the diagnostic accuracy of percutaneous Fistula contrast-enhanced ultrasound combined with transrectal 360o 3-D imaging and conventional transrectal ultrasound in complex anal Fistula. *Ultrasound Med Biol*. 2022;48(10):2154-2161.
<https://doi.org/10.1016/j.ultrasmedbio.2022.06.013>
417. Gui S, Patel N, Issenman R, et al. Acute management of pediatric cyclic vomiting syndrome: A systematic review. *J Pediatrics*. 2019;214:158-164. <https://doi.org/10.1016/j.jpeds.2019.06.057>
418. Eecke HV, Arnold D, Heyman S, et al. Cystoscope-guided trans-anal Fistula-tract Laser Closure (FiLaC) of late onset recurrent recto-urethral fistula post anorectal malformation repair: A case report. *J Pediatric Surgery Case Rep*. 2023;95:102674. <https://doi.org/10.1016/j.epsc.2023.102674>
419. Lin C-A, Chou C-M, Huang S-Y, et al. The optimal primary treatment for pediatric perianal abscess and anal fistula: A systemic review and meta-analysis. *J Pediatric Surgery*. 2023;58(7):1274-1280.
<https://doi.org/10.1016/j.jpedsurg.2023.01.055>
420. Bhusari S, Nikam K, Wakte P. Anal fistula: A comprehensive review. *Pharma Innov J*. 2019;8(1):201-208.
421. Santos CHMD, Cavasana AL, Moreira DDL, et al. Treatment of anal fistula with *Baccharis dracunculifolia* extract. Experimental study in rats. *Rev. Col. Bras Cir*. 2020, 46(5),e20192297.
<https://doi.org/10.1590/0100-6991e-20192297>
422. Comparin BC, Santos CHMD, Dourado DM, et al. Treatment of rats anal fistula with glycerin *Aloe barbadensis* Miller extract. *J. Coloproctol*. 2018, 38(4),283-289. <https://doi.org/10.1016/j.jcol.2018.05.009>
423. Mosleh G, Nimrouzi M, Badr P, et al. The approach of traditional Persian medicine to treatment of anal fissure. *Tradit. Integrat. Med*. 2019, 4(2), 78-83.
424. Cavasana AL, Santos CHMD, Dourado DM, et al. Effectiveness of the *Aloe vera* extract in the treatment of fistula-in-ano. *J. Coloproctol*. 2020, 40(1), 67-72.
<https://doi.org/10.1016/j.jcol.2019.09.003>
425. Zhang X, Liu W, Zhang S, et al. Wei-Tong-Xin ameliorates functional dyspepsia via inactivating TLR4/MyD88 by regulating gut microbial structure and metabolites. *Phytomedicine* 2022, 102, 154180.
<https://doi.org/10.1016/j.phymed.2022.154180>
426. Yang L, Liu X, Zhu J, et al. Progress in traditional Chinese medicine against chronic gastritis: From chronic non-atrophic gastritis to gastric precancerous lesions. *Heliyon* 2023, 9(6), e16754.
<https://doi.org/10.1016/j.heliyon.2023.e16754>
427. Schottenstein N, Yang D, Sanchez RE, et al. 455 efficacy and safety of aprepitant for children with cyclic vomiting syndrome and cannabinoid hyperemesis syndrome. *Gastroenterology* 2024, 166(5), S-100.
[https://doi.org/10.1016/S0016-5085\(24\)00725-X](https://doi.org/10.1016/S0016-5085(24)00725-X)
428. Laurindo LF, Maio MCD, Correa NDG, et al. Effects of medicinal plants and phytochemicals in Nrf2 pathways during inflammatory Bowel diseases and related colorectal cancer: A comprehensive review. *Metabolites*. 2023;13(2):243. <https://doi.org/10.3390/metabo13020243>
429. Abber SR, Randunne A, Staller K, et al. Su1647 psychological symptoms differentially predict symptoms between attacks and attack frequency in cyclic vomiting syndrome. *Gastroenterology*. 2024;166(5):S-780-S781.
[https://doi.org/10.1016/S0016-5085\(24\)02275-3](https://doi.org/10.1016/S0016-5085(24)02275-3)
430. Kilani Y, Egbo OJ, Markson F, et al. Mo1620 cannabis is associated with a reduced 30-day readmission of patients with cyclical vomiting syndrome. *Gastroenterology*. 2023;164(6):S-864.
[https://doi.org/10.1016/S0016-5085\(23\)02998-0](https://doi.org/10.1016/S0016-5085(23)02998-0)
431. Shah M, Jergel A, George RP, et al. Distinguishing clinical features of cannabinoid hyperemesis syndrome and cyclic vomiting syndrome: A retrospective cohort study. *J Pediatrics*. 2024;271:114054.
<https://doi.org/10.1016/j.jpeds.2024.114054>
432. Thavamani A, Umapathi KK, Khatana J, et al. Cyclic vomiting syndrome-related hospitalizations trends, comorbidities and health care costs in Children: A population based study. *Children*. 2022;9(1):55.
<https://doi.org/10.3390/children9010055>

433. Hayes WJ, VanGilder D, Berendse J, et al. Cyclic vomiting syndrome: diagnostic approach and current management strategies. *Clin Experim Gastroenterol.* 2018;2018:77-84. <https://doi.org/10.2147/CEG.S136420>
434. Kilani Y, Aljabiri Y, Arshad I, et al. Cannabis use and cyclical vomiting syndrome: An open debate. *Digest Liver Dis.* 2024;56(2):272-280. <https://doi.org/10.1016/j.dld.2023.10.002>
435. Bolender A, Staley R, Sclocco R, et al. Sa1615 abnormal temporal summation associated with primary somatosensory functional connectivity changes in response to pain challenge in cyclic vomiting syndrome. *Gastroenterology.* 2023;164(6):S-408. [https://doi.org/10.1016/S0016-5085\(23\)01943-1](https://doi.org/10.1016/S0016-5085(23)01943-1)
436. Ahmed O, Nedi T, Yimer EM. Evaluation of anti-gastric ulcer activity of aqueous and 80% methanol leaf extract of *Urtica simensis* in rats. *Metabol Open.* 2022;14:100172. <https://doi.org/10.1016/j.metop.2022.100172>
437. Akash SR, Tabassum A, Aditee LM, et al. Pharmacological insight of rutin as a potential candidate against peptic ulcer. *Biomed Pharmacother.* 2024;177:116961. <https://doi.org/10.1016/j.biopha.2024.116961>
438. Bashashati M, McCallum RW. Neurochemical mechanisms and pharmacologic strategies in managing nausea and vomiting related to cyclic vomiting syndrome and other gastrointestinal disorders. *Eur J Pharmacol.* 2014;722:79-94. <https://doi.org/10.1016/j.ejphar.2013.09.075>
439. Burow P, Haselier M, Naegel S, et al. The mitochondrial biomarkers FGF-21 and GDF-15 in patients with episodic and chronic migraine. *Cells.* 2021;10(9):2471. <https://doi.org/10.3390/cells10092471>
440. Chen C, Fu Y-H, Li M-H, et al. Nuclear magnetic resonance-based metabolomics approach to evaluate preventive and therapeutic effects of *Gastrodia elata* Blume on chronic atrophic gastritis. *J Pharmaceut Biomed Anal.* 2019;164:231-240. <https://doi.org/10.1016/j.jpba.2018.10.035>
441. Hikita T, Kodama H, Ogita K, et al. Cyclic vomiting syndrome in infants and children: A clinical follow-up study. *Pediatric Neurol.* 2016;57:29-33. <https://doi.org/10.1016/j.pediatrneurol.2016.01.001>
442. Kek H-P, Tsai W-L, Chiu P-C, et al. Navigating the diagnostic journey in pediatric gastroenterology: decoding recurrent vomiting and epigastric pain in a child with glutaric aciduria type II. *Children.* 2024;11(3):285. <https://doi.org/10.3390/children11030285>
443. Han C, Du D, Wen Y, et al. Chaqin chengqi decoction ameliorates acute pancreatitis in mice via inhibition of neuron activation-mediated acinar cell SP/NK1R signaling pathways. *J Ethnopharmacol.* 2021;274:114029. <https://doi.org/10.1016/j.jep.2021.114029>
444. Liao X, Tian Y, Zhang Y, et al. Acupuncture for functional dyspepsia: Bayesian meta-analysis. *Complement Ther Med.* 2024;82:103051. <https://doi.org/10.1016/j.ctim.2024.103051>
445. Mauritz MD, Hasan C, Schreiber L, et al. Differential diagnosis of cyclic vomiting and periodic headaches in a child with ventriculoperitoneal shunt: Case report of chronic shunt over-drainage. *Children.* 2022;9(3):432. <https://doi.org/10.3390/children9030432>
446. McAbee GN, Morse AM, Cook W, et al. Neurological etiologies and pathophysiology of cyclic vomiting syndrome. *Pediatric Neurol.* 2020;106:4-9. <https://doi.org/10.1016/j.pediatrneurol.2019.12.001>
447. Mehrabani S, Moslemi L. Management of resistant cyclic vomiting syndrome with a new empiric treatment in the prophylaxis period: A case report and review of literature. *J Pediatrics Rev.* 2021;9(1):67-70. <https://doi.org/10.32598/jpr.9.1.344.2>
448. Moses J, Keilman A, Worley S., et al. Approach to the diagnosis and treatment of cyclic vomiting syndrome: A large single-center experience with 106 patients. *Pediatric Neurol.* 2014;50(6):569-573. <https://doi.org/10.1016/j.pediatrneurol.2014.02.009>
449. Tarbell SE, Li BUK. Anxiety measures predict health-related quality of life in children and adolescents with cyclic vomiting syndrome. *J Pediatrics.* 2015;167(3):633-638. <https://doi.org/10.1016/j.jpeds.2015.05.032>
450. Tarbell SE, Millar A, Laudenslager M, et al. Anxiety and physiological responses to the Trier social stress test for children in adolescents with cyclic vomiting syndrome. *Autonomic Neurosci.* 2017, 202, 79-85. <https://doi.org/10.1016/j.autneu.2016.08.010>
451. Venkatesan T, Palsson OS, Iyer L, et al. Su1648 acute and long-term effects of heartfulness mediation on psychological outcomes and the endocannabinoid signaling system in cyclic vomiting syndrome using an APP. *Gastroenterology.* 2024;155(5):18-21. [https://doi.org/10.1016/S0016-5085\(24\)02276-5](https://doi.org/10.1016/S0016-5085(24)02276-5)
452. Wali R, Khan MF, Mahmood A, et al. Ethnomedicinal appraisal of plants used for the treatment of gastrointestinal complaints by tribal communities living in Diamir district, Western Himalayas, Pakistan. *Plos One.* 2022;17(6):e0269445. <https://doi.org/10.1371/journal.pone.0269445>
453. Song X, Chen YJ, Perry A, et al. Productivity loss and indirect burden of cyclic vomiting syndrome in the United States. *Gastro Hep Adv.* 2022;1(6):954-962. <https://doi.org/10.1016/j.gastha.2022.06.017>

454. Thavamani A, Sankararaman S. Cannabis use and cyclic vomiting syndrome. *Digest Liver Dis.* 2024;56(3):535. <https://doi.org/10.1016/j.dld.2023.11.018>
455. Trikola A, Bar N, Kiser K, et al. Mo1627 food-related symptoms and avoidance is prevalent even in between attacks in patients with severe cyclic vomiting syndrome. *Gastroenterology.* 2023;164(6):S-867. [https://doi.org/10.1016/S0016-5085\(23\)03005-6](https://doi.org/10.1016/S0016-5085(23)03005-6)
456. Venkatesan T, Hillard CJ, Rein L, et al. Patterns of cannabis use in patients with cyclic vomiting syndrome. *Clin Gastroenterol Hepatol.* 2020;18(5):1082-1090. <https://doi.org/10.1016/j.cgh.2019.07.039>
457. Chen YJ, Song X, Winer I, et al. Health care resource use and associated costs of cyclic vomiting syndrome in the United States. *Gastro Hep Adv.* 2022;1(6):963-973. <https://doi.org/10.1016/j.gastha.2022.06.013>
458. Clouse ER, Sayuk GS, Lustman PJ, et al. Zonisamide or levetiracetam for adults with cyclic vomiting syndrome: A case series. *Clin. Gastroenterol. Hepatol.* 2007;5(1):44-48. <https://doi.org/10.1016/j.cgh.2006.10.004>
459. Remes-Troche JM, Cerda ECM, Valdovinos-Garcia LR, et al. Su1652 clinical profile and associated factors in Mexican patients with cyclic vomiting syndrome/cannabinoid hyperemesis syndrome. *Gastroenterology.* 2024;166(5):S-782-S783. [https://doi.org/10.1016/S0016-5085\(24\)02280-7](https://doi.org/10.1016/S0016-5085(24)02280-7)
460. Rosman NP, Dutt M, Nguyen HT. A curable and probably often-overlooked cause of cyclic vomiting syndrome. *Seminars Pediatric Neurol.* 2014 21(2):60-65. <https://doi.org/10.1016/j.spen.2014.04.001>
461. Rubio-Tapia A, McCallum R, Camilleri M. AGA clinical practice update on diagnosis and management of cannabinoid hyperemesis syndrome: Commentary. *Gastroenterology.* 2024;166(5):930-934. <https://doi.org/10.1053/j.gastro.2024.01.040>
462. Shahrajabian MH. Medicinal plants related to antidiabetic properties from Latin America. *J. Stress Physiol. Biochem.* 2025;21(4):215-245.